

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Fungicide



CERATO 250 EC

Reg. No. L 9336 Act/Wet No. 36 of/van 1947

1:19/8/2013-Nov2018

*An emulsifiable concentrate contact and translaminar fungicide for the control of Black spot in citrus and Black shank on flue-cured tobacco.
'n Emulgeerbare konsentraat kontak- en translaminêre swamdoder vir die beheer van Swartvlek in sitrus en Swartstam op oongedroogde tabak.*

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

pyraclostrobin (strobilurin) **250 g/l** piraklostrobien (strobilurien)

FRAC FUNGICIDE GROUP CODE **11** FRAC SWAMDODER GROEPKODE



villa 

Registration holder / Registrasiehouer:

Villa Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1992/002474/07

PO Box / Posbus 10413, Aston Manor, 1630

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: 2902

Willow Set & Print 011 394-4486



HARMFUL
SKADELIK



GEBRUIKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED
REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotonommer:

CERATO 250 EC

Reg. No. L 9336 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE /FRAC SWAMDODER GROEPKODE: 11

ACTIVE INGREDIENT/ AKTIEWE BESTANDDEEL:

pyraclostrobin (strobilurin) / piraklostrobien (strobilurien).....250 g/l

Registration holder / Registrasiehouer:

VILLA CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1992/002474/07 Mpy. Reg. Nr.

P.O. Box / Posbus 10413, ASTON MANOR, 1630

Tel. (011) 396 2233

HARMFUL  **SKADELIK**

WARNINGS**Withholding periods:**

Minimum number of days between the last application and harvest:	
Citrus: Oranges, Grape fruit and Lemons	45 days
Tobacco	83 days

IN THE CASE OF CROPS INTENDED FOR EXPORT, CONFIRM WITHHOLDING PERIODS WITH THE RELEVANT EXPORT ORGANISATION. IF THIS PRODUCT IS USED ACCORDING TO THE RECOMMENDATIONS ON THIS LABEL, LOCAL MRL VALUES WILL NOT BE EXCEEDED. HOWEVER, VILLA CROP PROTECTION (PTY) LTD. CANNOT ACCEPT RESPONSIBILITY FOR EXPORTED CROPS EXCEEDING THE IMPORT TOLERANCES OF OTHER COUNTRIES.

- Handle with care.
- Harmful when swallowed.
- Store in a cool, dry place away from food, feeds, fertilizers and other agricultural remedies.
- Irritating to eyes and skin.
- Toxic to fish and aquatic invertebrates.
- Keep out of reach of children, animals and uninformed persons.
- Re-entry: Do not enter treated area for 1 day after application unless wearing protective clothing.
- **In case of poisoning, call a doctor and make this label available to him / her.**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Avoid eye and skin contact and inhalation of spray mist.
- Wear rubber gloves and a face shield during mixing and application.
- Do not enter treated areas until the sprayed solution has dried.
- Wash contaminated clothing after use.
- Wash with soap and water after use or accidental skin contact.
- Do not eat, drink or smoke whilst mixing, applying the product or before washing hands and face or change of clothing.
- Avoid drift of spray onto other crops, grazing, rivers, dams and areas not under treatment.
- Clean the applicator with a 1 % household ammonia solution before using for other pesticides. Let the solution stand for several hours, preferably overnight, in the spray equipment. Rinse at least twice. This applicator should not be used for applying chemicals other than pesticides.

- Thoroughly clean the spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- Containers and packages must be completely emptied before being disposed of. Shake out thoroughly into the applicator and destroy the empty container thereafter in the described manner.
- Invert the empty container over the spray or mixing tank to make sure the container is empty before destroying the container.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of a third of the volume of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening.
- In case of spillage or leakage, soak up with soil or other absorbent materials and dispose of at any approved waste disposal facility.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

FIRST AID TREATMENT

- Remove the patient from the source of poisoning and keep him/her calm and at rest. **Get medical attention immediately.**
- **Inhalation:** Remove the patient from the exposure area to fresh air immediately. Keep affected person calm and reassured. Get medical attention immediately.
- **Skin contact:** Immediately remove contaminated clothing, shoes and leather goods (e.g. watch bands and belts). Wash the contaminated skin gently and thoroughly with non-abrasive soap and water. **Get medical attention immediately.**
- **Eye contact:** Immediately flush eyes with copious quantities of lukewarm, gently flowing water for at least 20 minutes or until the product is removed, while holding the eyelid(s) open. Take care not to rinse contaminated water into the unaffected eye or onto the face. If irritation, redness or burning sensations develop, **get medical attention immediately.**
- **Ingestion:** **Seek medical advice immediately** and show this label.

NOTE TO PHYSICIAN

Treat according to symptoms (decontamination, vital functions), no known specific antidote.

RESISTANCE WARNING

CERATO 250 EC is a group code 11 fungicide. Any fungus population may contain individuals naturally resistant to **CERATO 250 EC** and other group code 11 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly and exclusively in programs. These resistant fungi may not be controlled by **CERATO 250 EC** or any other group code 11 fungicides.

To delay fungicide resistance:

- avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate with products from different fungicide group codes,
- integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management contact the registration holder of this product.

USE RESTRICTIONS

- **CERATO 250 EC** must not be used on old neglected citrus trees, as they are extremely prone to fungicidal resistance.
- Do not apply **CERATO 250 EC** on non-tolerant Black shank tobacco cultivars.
- **CERATO 250 EC** may only be applied on flue cured tobacco.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

- **CERATO 250 EC** is compatible with **Multizeb 455 SC, Villa Unizeb 800 WP / Cozeb 800 WP** and **Copper oxychloride WP** and light to medium narrow range mineral oils.
- If tank mixtures with other products are made, first confirm compatibility by mixing small volumes of the products in the correct ratio with the appropriate quantity of water.
- Water quality and formulation properties of other products may influence compatibility.

Mixing instructions:

- Half fill the spray tank with clean water and ensure agitation.
- Pre-mix the required amount of **Mancozeb** or **Copper** containing product, in at least 10 litres of water and add it to the spray tank.
- Pre-mix the required volume of **CERATO 250 EC** in at least 10 litres of water before adding to the spray tank.
- Citrus only: Finally add the mineral oil to the mixture. Fill the spray tank with water to the final volume required while maintaining agitation, to ensure thorough mixing of the spray mixture before spraying commences.
- Maintain agitation while spraying.
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

CROP RECOMMENDATIONS**Citrus:**

CERATO 250 EC must always be applied in a programme application with a **Mancozeb** or a **Copper** containing product in order to achieve a responsible resistance management programme. Accurate timing of treatments is essential to achieve efficient control.

- **CERATO 250 EC** treatments must be applied as a preventive treatment with limited curative abilities.
- Apply a **Mancozeb** or a **Copper** containing product as a first treatment starting at fruit set (100 % petal drop), at the registered rate.
- Follow up with two (2) **CERATO 250 EC** applications, as recommended below. Complete the spray programme with a further treatment of at least one **Mancozeb** or **Copper** containing product application, as recommended.
- Refer to the respective **Mancozeb**, **Copper** containing products and narrow range mineral oil product labels for the respective recommendations and application rates.
- **Lemons for export crop and local oil production:** All bearing and out of season fruit, should be removed from the trees before starting the programme application. Do not deviate from the recommended spray programme starting at fruit set (100 % petal drop).

Tobacco:

- Apply **CERATO 250 EC** in a spray programme in order to optimise the fungicidal efficacy of the product and to ensure sustainable use of strobilurins. Apply the programme as follows:
 - first application of **CERATO 250 EC** at transplanting,
 - follow up four (4) weeks later with **Vimethozeb 690 WP**.
- Only apply **CERATO 250 EC** on flue cured tobacco.

Application instructions:

All spray operations must be performed with suitable equipment that is in good working order and correctly calibrated to give the desired coverage for that particular method of application.

APPLICATION RATES

Crop / Disease	Dosage rate	Remarks
<u>Tobacco</u> Black shank (<i>Phytophthora nicotianae</i>)	1.2 l CERATO 250 EC per 15000 plants	Refer to the “ CROP RECOMMENDATIONS ” above. <u>Application at transplanting:</u> Apply before transplanting of tobacco. Dilute 80 ml CERATO 250 EC in 100 litres water. Fill the planting hole with water and apply 100 ml of the mixture into the planting hole while still full with water. <u>Application 4 weeks after transplant:</u> Follow the CERATO 250 EC treatment up 28 days later with Vimethozeb 690 WP as per relevant label instructions.

Crop / Disease	Dosage rate	Remarks
Citrus Oranges, Grape fruit & Lemons Black spot (<i>Guignardia citricarpa</i>)	200 g / 100 ℓ water Villa Unizeb 800 WP / Cozeb 800 WP OR 350 mℓ Multizeb 455 SC	Refer to “ CROP RECOMMENDATIONS ” above. Apply one (1) Mancozeb application at fruit set (100 % petal drop).
	10 mℓ CERATO 250 EC PLUS 300 mℓ light to medium narrow range mineral oil / 100 ℓ water PLUS 150 g / 100 ℓ water Villa Unizeb 800 WP / Cozeb 800 WP OR 260 mℓ / 100 ℓ water Multizeb 455 SC	Apply CERATO 250 EC 21 to 25 days after the first Mancozeb application as a preventive treatment. Ensure good coverage of all fruit with a medium to full cover spray. Apply a second CERATO 250 EC application 6 weeks later.
	200 g / 100 ℓ water Villa Unizeb 800 WP/ Cozeb 800 WP OR 350 mℓ / 100 ℓ water Multizeb 455 SC	Apply Mancozeb six (6) weeks after the second CERATO 250 EC application. Repeat with a Mancozeb treatment 21 to 24 days later if an infection period occurs in February.
Citrus: REMARKS Copper oxychloride WP product application as an alternative to Mancozeb in a CERATO 250 EC spray programme on citrus. <ul style="list-style-type: none"> As an alternative to Mancozeb (solo or in combination with CERATO 250 EC and mineral oil), Copper oxychloride WP can be included in the programme application. The Copper oxychloride WP applications should be applied according to the registered rates (Refer to the necessary label recommendations). In order to avoid/reduce the occurrence of Copper rind stippling and any lesion accentuation on fruit, Copper oxychloride WP should be applied at least 60 days apart. Reduce the number of Copper oxychloride WP applications to a maximum of 2 to 3 applications per season. 		

The following products mentioned in this label can be replaced with the following equivalent products:

- VILLA UNIZEB 800 WP** (L 8056 / N-AR 1091) = **COZEB 800 WP** (L 7098) = **NEWZEB 800 WP** (L 8167) (**Mancozeb**),
- MULTIZEB 455 SC** (L 9174) = **NEWZEB 455 SC** (L 9175) (**Mancozeb**),
- COPPER OXYCHLORIDE WP** (L 6231) = **COPPER OXYCHLORIDE WP** (L 5809) and

- **VIMETHOZEB 690 WP** (L 7986 / N-AR 1296) = **DIMETHOZEB 690 WP** (L 7985) (**Dimetomorph + Mancozeb**).

COZEB 800 WP, NEWZEB 455 SC, DIMETHOZEB 690 WP and/en **COPPER OXYCHLORIDE WP** are registered products of / is geregisteerde produkte van **UNIVERSAL CROP PROTECTION (PTY) LTD.**

VILLA UNIZEB 800 WP, MULTIZEB 455 SC, VIMETHOZEB 690 WP and/en **COPPER OXYCHLORIDE WP** are registered products of / is geregisteerde produkte van **VILLA CROP PROTECTION (PTY) LTD.**

NEWZEB 800 WP is a registered product of / is 'n geregisteerde produk van **CROPASURE (PTY) LTD.**