

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Adjuvant



GLYPHO-TOP

Reg. No. L 9604 Act/Wet No. 36 of/van 1947

1:22/7/2014 – May 2017

An aqueous water conditioning agent that will reduce alkaline water pH, complex certain antagonistic ions in carrier water and enhance the activity and compatibility of Glyphosate.

'n Vloeibare oplossing wat alkaliese spuitwater pH verlaag, sekere antagonistiese ione in die spuitwater sal bind, en die aktiwiteit en mengbaarheid van Glifosaat sal verbeter.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

ammonium sulphate 250 g/l ammoniumsulfaat



villa

Registration holder / Registrasiehouer:

Villa Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1992/002474/07

PO Box / Posbus 10413, Aston Manor, 1630

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: Not Regulated

Willow Set & Print 011 394-4486



**CAUTION
VERSIGTIG**



GEBRUIKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

DIRECTIONS FOR USE ENCLOSED

Batch number:
Lotnommer:

GLYPHO-TOP

Reg. No. L 9604 Act/Wet No. 36 of/van 1947

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

ammonium sulphate / ammoniumsulfaat250 g/l

Registration holder / Registrasiehouer:

VILLA CROP PROTECTION (PTY) LTD

PO Box / Posbus 10413

ASTON MANOR, 1630

Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS**

- Adhere to all label warnings that apply to the product or products with which **GLYPHO-TOP** is to be used.
- May irritate the eyes and skin.
- Store away from food, feeds, seed, fertilizers and other agricultural chemicals.
- Keep out of reach of children, uninformed persons and animals.
- Store in a cool, dry, well-ventilated place.
- Re-entry: Adhere to the re-entry interval and requirements as indicated on the herbicide label.

Aerial application:

Notify all the inhabitants in the immediate vicinity of the area to be sprayed and issue the necessary warnings.

Although GLYPHO-TOP has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weeds against the herbicide concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Avoid inhalation of the spray mist.
- Wear protective clothing: overalls, rubber boots and gloves, when handling the product.
- Wash thoroughly with soap and water after use and after accidental skin contact.
- In the case of eye contact flush eyes with copious amounts of water for at least 15 minutes.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and change of clothing.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams or areas not under treatment.
- Thoroughly clean the applicator after use and dispose of wash water in a way that will not contaminate food, grazing, boreholes, rivers and dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening and dispose of it in a safe way.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

PRODUCT PROPERTIES

- **Glyphosate** requires the pH of spray solution to be slightly acidic (normally pH 4.5 to 5.5). **Glyphosate** can also be antagonized by ions dissolved in carrier water. **GLYPHO-TOP** in combination with **Glyphosate** will decrease the pH of the spray solution and will complex antagonistic ions such as calcium, magnesium, sodium and potassium. The spray solution pH will, however, not be decreased to the extent that incompatibility with phenoxy herbicides is experienced.
- Additional acidification with other buffers and/or acidifiers is not recommended.
- The use of **GLYPHO-TOP** will increase **Glyphosate** efficacy by improving its uptake and may aid compatibility of **Glyphosate** with certain other herbicides.

USE RESTRICTIONS

- **GLYPHO-TOP** should not be used with any herbicide not listed on this label as it may improve the efficacy of certain herbicides to such an extent that it may cause damage to the crop.
- Take all necessary precautions associated with aerial spraying of herbicides especially with regard to control of drift.
- Test the mixture on a small scale before applying it on a large scale.

DIRECTIONS FOR USE: Use only as directed.**Compatibility:**

- **Scat[®] 360 SL, Mallet 100 SL, Imazethapyr 100 SL, Panga 360 SL, Slash 360 SL, Slash 710 SG, Glygran 710 SG, Slash Turbo 450 SL, Panga Turbo 450 SL, Slash Plus 540 SL, Panga Plus 540 SL, Villa 51.**
- Consult the labels of the abovementioned products for full particulars.

Mixing Instructions:

- Half fill the spray tank with clean water.
- Add the required amount of **GLYPHO-TOP** followed by the herbicide and then the remainder of the water while stirring.
- In the event of tank mixes, add **GLYPHO-TOP** first, then the complementary herbicide followed by the correct amount of the foliar active herbicide and then the remaining water with continual agitation.
- Should a wetter/spreader be required, add this last and only use **Villa 51** or an approved Villa Crop Protection surfactant.
- Maintain adequate agitation during application and the prepared mixture must not be left in the spray tank for any length of time.
- **GLYPHO-TOP** is corrosive; therefore the sprayer, tank and system should be rinsed thoroughly after the use of **GLYPHO-TOP**, to remove all ammonium sulphate residues.

Application Instructions:**Ground application:**

GLYPHO-TOP can be used with any correctly calibrated spray equipment as recommended on the herbicide label.

Aerial application:

Take all necessary precautions associated with aerial spraying of herbicides especially with regard to control of drift. Follow the label instructions of the herbicide being used with **GLYPHO-TOP**. If in any doubt, consult with the supplier and aerial crop sprayer.

APPLICATION RATES

- If possible, test the Electrical Conductivity or have an analysis of the spray water done. The precise rate of **GLYPHO-TOP** can then be determined by your Villa Crop Protection representative. The analysis report must contain values for calcium, magnesium, sodium and potassium content **OR** an Electrical Conductivity measurement when a full analysis is not available. If this is not possible, use the dosage rate table as a guideline.
- In cases where a water analysis or an Electrical Conductivity measurement is not available, add 0.5 to 2.0 litres **GLYPHO-TOP** to 100 litres clean water (preferably not murky) to obtain a 0.5 to 2.0 % spray solution. If in any doubt about the spray water quality/Electrical Conductivity, use the higher recommended **GLYPHO-TOP** rate.

DOSAGE RATE OF GLYPHO-TOP IF A WATER ANALYSIS IS NOT AVAILABLE OR IF A VILLA REPRESENTATIVE HAS NOT MADE A PRECISE CALCULATION				
Electrical Conductivity (EC)			GLYPHO-TOP DOSAGE RATE	
Milli Siemens per m (mS/m)	Milli Siemens per cm (mS/cm)	Micro Siemens per cm (μS/cm)	%	Litres per 100 litres water
0 to 100	0 to 1.0	0 to 1000	0.5 to 1.0 %	0.5 to 1.0 litres per 100 litres water
101 to 200	1.1 to 2.0	1001 to 2000	1.0 to 2.0 %	1.0 to 2.0 litres per 100 litres water
> 200	> 2.0	> 2000	2.0 %	2.0 litres per 100 litres water

NOTE

If a water analysis is available that indicates the levels of calcium (Ca²⁺), magnesium (Mg²⁺), sodium (Na⁺) and potassium (K⁺) **OR** Electrical Conductivity, your **Villa Crop Protection** representative can determine the precise amount of product to be used. This calculated amount may be outside the 0.5 to 2 % rate range, depending on the quality of spray water.

The following product mentioned in this label may be replaced with an equivalent product:

- **VILLA 51** (L 8050 / W 130454 / N-AR 1090) = **WEN 51** (L 8315).
- **SCAT® 360 SL** (L 5716 / W 130453) = **SLASH 360 SL** (L 7811 / N-AR 1087) = **PANGA 360 SL** (L 7812) = **GLYPHOSATE 360 SL** (L8128),
- **MALLET 100 SL** (L 8049) = **IMAZETHAPYR 100 SL** (L 8051),
- **SLASH 710 SG** (L 8450 / W 130058) = **GLYGRAN 710 SG** (L 8449) = **LYNCH 710 SG** (L8670),
- **SLASH TURBO 450 SL** (L 8505) = **PANGA TURBO 450 SL** (L 8506) = **LYNCH TURBO 450 SL** (L8754),
- **SLASH PLUS 540 SL** (L 8819) = **PANGA PLUS 540 SL** (L 8818) = **LYNCH PLUS 540 SL** (L8816).

SLASH 360 SL, IMAZETHAPYR 100 SL, SLASH 710 SG, SLASH TURBO 450 SL and/en SLASH PLUS 540 SL are registered products of / is geregistreerde produkte van **Universal Crop Protection (Pty) Ltd.**

SCAT® 360 SL is a registered trademark of / is 'n geregistreerde handelsmerk van **Universal Crop Protection (Pty) Ltd.**

PANGA 360 SL, GLYGRAN 710 SG, PANGA TURBO 450 SL, PANGA PLUS 540 SL and/en MALLET 100 SL are registered products of / is geregistreerde produkte van **Villa Crop Protection (Pty) Ltd.**

GLYPHOSATE 360 SL, LYNCH 710 SG, LYNCH TURBO 450 SL and/en LYNCH PLUS 540 SL are registered products of / is 'n geregistreerde handelsmerk van **Crop Asure (Pty) Ltd.**