

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Fungicide



SCOPE 250 EC

Reg. No. L 7942 Act/Wet No. 36 of/van 1947  
N-AR 1098

3: 10/3/2016-Aug2019

*An emulsifiable concentrate fungicide with systemic action for the control of diseases on crops as listed.*

*'n Emulgeerbare konsentraat swamdoder met sistemiese werking vir die beheer van siektes op gewasse soos aangedui.*

**ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL**  
difenoconazole (triazole) **250 g/l** difenokonasool (triasool)

FRAC FUNGICIDE GROUP CODE **3** FRAC SWAMDODER GROEPKODE



**villa** 

Registration holder / Registrasiehouer:

Villa Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1992/002474/07

PO Box / Posbus 10413, Aston Manor, 1630

Tel: 011 396 2233

Website / Webblad: [www.villacrop.co.za](http://www.villacrop.co.za)

UN Number: 3082

Willow Set & Print 011 394-4486



CAUTION  
VERSIGTIG



GEbruiksAANwYsINGS INGESLUIT

VERWYS NA BESONDERHEDE  
GEDRUK OP HOUER/SAK

Date formulated:

Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED  
ON CONTAINER/BAG

Batch number:

Lotnummer:



**SCOPE 250 EC**

Reg. No. L 7942 / N-AR 1098 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: 3

**ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:**

difenoconazole (triazole) / difenokonasool (triasool) ..... 250 g/l

Registration holder / Registrasiehouer:

**VILLA CROP PROTECTION (PTY) LTD.**

Co. Reg. No. 1992/002474/07 Mpy. Reg. Nr.

PO Box / Posbus 10413

ASTON MANOR, 1630, Tel. (011) 396 2233

**CAUTION / VERSIGTIG****WARNINGS****Withholding periods:**

| <b>Minimum time between last application and harvest or feeding:</b> |         |
|--|---------|
| Apples and Pears   | 21 days |
| Citrus   | 90 days |
| Beans  | 14 days |
| Grapes   | 28 days |
| Groundnuts (fodder)  | 21 days |
| Maize (do not graze or use treated maize as fodder)                  | 14 days |
| Soybeans   | 18 days |
| Potatoes   | 14 days |
| Tomatoes   | 7 days  |

**NOTE**

**THE RECOMMENDED WITHHOLDING PERIODS MEET LOCAL RESIDUE LEVEL REQUIREMENTS AND NOT NECESSARILY THE REQUIREMENTS FOR EXPORT. WHEN THE CROP IS DESTINED FOR EXPORT, CONSULT THE REQUIREMENTS AND GUIDELINES FOR EXPORT BEFORE THE PRODUCT IS USED.**

- Handle with care.
- Harmful if swallowed, inhaled or when absorbed through the skin.
- Risk of serious damage to eyes.
- Mild skin irritant.
- Toxic to fish and aquatic organisms.
- Avoid storage temperatures below 10 °C and above 35 °C.
- Store in the original container in a cool, dry, well-ventilated place away from food, seed and feedstuffs, fertilizers and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

**Aerial application:**

Notify all inhabitants in the immediate vicinity of the area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate adjacent areas or water sources.

**Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.**

**PRECAUTIONS**

- Avoid inhalation of the spray mist.
- Avoid eye and skin contact.
- Wear a face shield and rubber gloves when handling the product, preparing the spray mixture and during application.
- Wash with soap and water immediately after use and accidental skin contact.
- Wash contaminated clothing after use.
- In case of eye contamination, rinse the eyes thoroughly with plenty of clean, water for at least 20 minutes, holding the eyelid(s) open. If irritation persists obtain medical assistance immediately.
- Do not eat, drink or smoke whilst mixing and applying the product or before washing hands and face and change of clothing.
- Prevent drift onto other crops, grazing, rivers, dams, boreholes and areas not under treatment or to nearby water sources.
- Thoroughly clean spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of a third of the volume of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the container by perforation and flattening and dispose of it in a safe way.
- **Do not** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

**RESISTANCE WARNING**

**SCOPE 250 EC** is a group code 3 fungicide. Any fungus population may contain individuals naturally resistant to **SCOPE 250 EC** and any other group code 3 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **SCOPE 250 EC** or any other group code 3 fungicide.

To delay fungicide resistance:

- avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes,
- for tank mixing or alternation with products in fungicide group code M, refer to applicable individual product labels,
- integrate other control methods (chemical, cultural, biological) into disease control programmes.

For specific information on resistance management contact the registration holder of this product.

**USE RESTRICTIONS**

- The uptake and activity of systemic compounds may be adversely affected when crops experience water related stress and/or fertility stress conditions. It is therefore not advised to apply **SCOPE 250 EC** during periods of stress. If in doubt, consult Villa Crop Protection (Pty) Ltd. or a distributor.
- Spray programmes with **SCOPE 250 EC** should be preventative unless mentioned otherwise.
- **As an anti-resistance strategy, it is not recommended to apply SCOPE 250 EC (or related fungicides) exclusively for prolonged periods during the season.**

**DIRECTIONS FOR USE: Only use as indicated.****General properties:**

- **SCOPE 250 EC** is a local systemic fungicide with limited acropetal movement.
- If applied in a protective spray programme **SCOPE 250 EC** effectively controls the following diseases: **Brown rust** (*Uromyces appendiculatus*) in beans, **Early leafspot** (*Cercospora arachidicola*), **Late leafspot** (*Cercosporidium personatum*), **Web blotch** (*Phoma arachidicola*) and **Rust** (*Puccinia arachidis*) in groundnuts, **Early blight** (*Alternaria solani*) in potatoes and tomatoes, **Grey leafspot** (*Cercospora zea-maydis*) and **Northern leaf blight** (*Helminthosporium turcicum*) in maize and sweetcorn, **Apple scab** (*Venturia inaequalis*) in apples, **Pear scab** (*V. pirina*) in pears, **Powdery mildew** (*Oidium tuckerii*) in grapes, **Soybean rust** (*Phakopsora pachyrhizi*) on soybeans and **Navel-end-rot / Black core rot** (*Alternaria citri*) and **Brown spot** (*A. alternata*) in citrus.

**Compatibility:**

- **SCOPE 250 EC** is compatible with the following fungicides, **Mycoblock 250 SC**, **Inspect 400 SC**, **Villa Unizeb 800 WP** and **Captab 500 SC** as indicated in the label recommendations. The compatibility of **SCOPE 250 EC** in tank mixtures with other products has not been confirmed and can be influenced by the formulations involved, as well as the water quality.
- The compatibility of **SCOPE 250 EC** with other products may vary. A physical compatibility test must always be performed before such tank mixture is sprayed.
- When **SCOPE 250 EC** is used in conjunction with any other agricultural remedy, all **WARNINGS**, **PRECAUTIONS** and **DIRECTIONS FOR USE** mentioned on that label, must be adhered to.

**Mixing instructions:**

- Half fill the spray tank with clean water.
- Shake the **SCOPE 250 EC** container well before use.
- Add the required quantity of **SCOPE 250 EC** to the water in the spray tank.
- Agitate well to obtain a homogeneous emulsion.
- Fill the spray tank with water to the required level while maintaining agitation, to ensure thorough mixing.
- Maintain agitation while spraying.
- If a wettable powder is to be sprayed in a mixture with the **SCOPE 250 EC**:
  - a) pre-cream the wettable powder with clean water,
  - b) add the pre-creamed wettable powder into the spray tank, while ensuring agitation,
  - c) then add the **SCOPE 250 EC** and
  - d) fill the spray tank with water to the required level.
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

**APPLICATION****Ground application:**

- **SCOPE 250 EC** can be applied with conventional high volume spray equipment.
- It is important that the sprayer be calibrated prior to application to ensure that the correct delivery rate and even distribution of the spray mixture is obtained.
- Apply in 300 to 500 litres water per hectare.
- It is recommended that the nozzles should be fitted with hollow cone tips giving medium to fine droplets.

**Aerial application:**

Aerial application of **SCOPE 250 EC** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- Volume: A spray mixture volume of 30 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.
- Droplet coverage: 25 to 35 droplets per cm<sup>2</sup> must be recovered at the target area.
- Droplet size: A droplet spectrum with a VMD of 280 to 300 micron is recommended. Limit the production of fine droplets, less than 150 micron (high drift and evaporation potential), to a minimum.
- Flying height: Maintain the height of the spray boom at 3 to 4 meters above the target. Do not spray when aircraft is diving, climbing or when banking.
- Use suitable atomising equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8°C.
- Stop spraying if the wind speed exceeds 15 km per hour.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:

- a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),
- b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

**APPLICATION RATES**

| <b>Crop &amp; Disease</b>   | <b>Dosage rate</b>   | <b>Remarks</b>        |
|---|--|-----------------------|
| <b>Apples and Pears</b><br>Scab<br>( <i>Venturia</i> spp)   | Apply <b>SCOPE 250 EC</b> in a programmed spray schedule as a full cover spray to point of run-off or in a concentrated form (4x).           |                       |
| <b>Development stage</b>  | <b>High volume dosage rate / 100 ℓ water</b>   | <b>Spray interval</b> |
| From green tip and throughout the pre-blossom period  | 6 ml <b>SCOPE 250EC</b><br><b>PLUS</b><br>100 ml <b>Captab 500 SC</b><br><b>OR</b><br>Villa Unizeb 800 WP<br>at a rate of 80 g / 100 ℓ water | 7 days                |
| During post-blossom period  | 6 ml <b>SCOPE 250EC</b><br><b>PLUS</b><br>100 ml <b>Captab 500 SC</b><br><b>OR</b><br>Villa Unizeb 800 WP<br>at a rate of 80 g / 100 ℓ water | 10 to 14 days         |
| <b>GENERAL REMARKS</b>  |  |                       |
| <ul style="list-style-type: none"> <li>Consult the <b>Inspect 400 SC</b>, <b>Captab 500 SC</b> and <b>Villa Unizeb 800 WP</b> labels for <b>USE RESTRICTIONS</b>, <b>WARNINGS</b> (including <b>RESISTANCE WARNINGS</b> and <b>WITHHOLDING PERIODS</b>), <b>PRECAUTIONS</b> and <b>DIRECTIONS FOR USE</b>.</li> <li><b>SCOPE 250 EC</b> is recommended in a protective spray programme in areas where and during periods when conditions are favorable for the development of Scab. During December it is recommended to change to a protective <b>contact fungicide spray programme</b>.</li> <li>When a protective <b>contact fungicide spray programme</b> is followed, <b>SCOPE 250 EC</b> should be applied as a curative treatment within two (2) days of the start of an infection period. In such situations use 10 ml <b>SCOPE 250 EC PLUS Captab 500 SC OR Villa Unizeb 800 WP</b> at the above indicated rates per 100 litres water <b>OR</b> with 56 ml <b>Inspect 400 SC</b> per 100 litres water. Curative applications are only recommended when adequate and reliable disease prediction systems are followed.</li> </ul> |  |                       |
| <b>NOTES REGARDING APPLICATION ON APPLES AND PEARS</b>  |  |                       |
| <b>High volume application (HV):</b>  |  |                       |
| <ul style="list-style-type: none"> <li>The water volume to be applied per hectare must be calculated according to the Tree Row Volume (TRV) formula:</li> </ul>   |  |                       |
| $\text{HV } \ell/\text{ha} = \frac{\text{Tree height} \times \text{Tree diameter} \times 937}{\text{Row width}}$  |  |                       |
| <ol style="list-style-type: none"> <li>This is the water volume required when the trees are in full leaf.</li> <li>Use the water volume required per hectare to calculate the amount of <b>SCOPE 250 EC</b> required per hectare.</li> <li>The following water requirements per hectare are recommended according to the different growth stages of the tree:</li> </ol>  |  |                       |
| <b>Growth stage</b>   | <b>% Water volume per hectare with regard to high volume (HV calculated above)</b>   |                       |
| From green tip to ± 30 % blossom  | 60 %   |                       |
| From ± full blossom to ± mid-November   | 80 %   |                       |
| From ± beginning December   | 100 %  |                       |
| <ul style="list-style-type: none"> <li><b>Low Volume (4x):</b><br/>When <b>SCOPE 250 EC</b> is applied as a low volume spray, ensure that the correct amount of product is applied per hectare.</li> </ul>  |  |                       |

| <b>Crop &amp; Disease</b>  | <b>Dosage rate</b>   | <b>Remarks</b>   |
|--|--|--|
| <b>Beans</b><br>Brown rust<br>( <i>U. appendiculatus</i> )   | Add a Villa approved oil adjuvant to <b>SCOPE 250 EC</b> . Apply in a protective spray programme, commencing at the start of flowering or sooner on highly susceptible cultivars, or at the first signs of disease. Treatments should be applied during periods of active plant growth and the programme should continue until plants start to senescence. |  |
|  | 250 ml / ha<br><b>OR</b><br>325 ml / ha  | <b>Ground application:</b><br>Apply the lower dosage every 10 to 14 days or the higher dosage every 18 to 21 days in the case of dry beans.  |
|  | 300 ml / ha  | <b>Aerial application:</b><br>Apply every 10 to 14 days (dry beans).   |
| <b>Citrus</b><br>(Navels, Clementine and Mandarin)<br>Navel-end-rot / Black core rot ( <i>A. citri</i> )   | 40 to 60 ml / 100 l water  | Apply twice. The first application at 50 % petal drop and the second at 100 % petal drop (when all the petals on the trees in the orchard have dropped). Apply as a medium cover spray ( $\pm$ 20 litres spray volume per tree, depending on tree size).   |
| <b>Mandarin</b><br>Brown Spot<br>( <i>A. alternata</i> )   | 40 to 60 ml / 100 l water<br><b>PLUS</b><br>200 g / 100 l water<br><b>Villa Unizeb 800 WP</b>  | Use the higher rate on orchards having a history of <i>Alternaria</i> diseases.  |
| <b>NOTE</b>  |  |  |
| <ul style="list-style-type: none"> <li>Do not apply after 100 % petal fall.</li> <li>Do not apply more than 3.0 litres per hectare <b>SCOPE 250 EC</b> per season.</li> <li>The two <b>SCOPE 250 EC</b> applications are part of a season-long control strategy for <i>Alternaria alternata</i> (Brown spot) control.</li> </ul> |  |  |
| <b>Groundnuts</b><br>Early leafspot;<br>( <i>C. arachidicola</i> )<br><br>Late leafspot;<br>( <i>C. personatum</i> )<br><br>Web blotch &<br>( <i>P. arachidicola</i> )<br><br>Rust<br>( <i>P. arachidis</i> )  | 200 ml and 300ml / ha<br><b>OR</b><br>325 ml / ha  | <b>Ground application: - Dry land</b><br>Add a Villa approved oil adjuvant to <b>SCOPE 250 EC</b> . Commence application at the first signs of disease. Use 300 ml per hectare for the first application and repeat every 10 to 14 days using 200 ml per hectare. Alternatively, apply the 325 ml per hectare dosage rate every 18 to 21 days. |
|  | 300 ml / ha<br><b>OR</b><br>400 ml / ha  | <b>Ground application: – Irrigation</b><br>Apply in a protective spray programme, commencing at the start of flowering (10 to 12 weeks after planting) or earlier, when the disease(s) starts to develop. Apply the lower dosage every 10 to 14 days or the higher dosage every 18 to 21 days.   |
|  | 350 ml / ha  | <b>Aerial application:</b><br>Aerial application is an emergency measure when conditions are unfavourable for ground application. Apply every 2 to 3 weeks on dryland crops and every 10 to 14 days on irrigated crops. All spray programmes should continue until the plants start to senescence naturally.                                   |

| <b>Crop &amp; Disease</b>   | <b>Dosage rate</b>   | <b>Remarks</b>   |
|---|--|--|
| <p><b>Maize and Sweetcorn</b><br/>Grey leaf spot<br/>(<i>Cercospora zeae-maydis</i>)</p> <p>Northern leaf blight<br/>(<i>Exserohilum turcicum</i> = <i>Helminthosporium turcicum</i>)</p> | <p><b>Broadcast Application (Ground &amp; Aerial):</b><br/>Apply <b>SCOPE 250 EC</b> as part of a spray programme with <b>Mycoblock 250 SC</b> as follows (refer to the <b>Mycoblock 250 SC</b> label):</p> <ul style="list-style-type: none"> <li>Northern leaf blight - apply the first application during the 8- to 10-leaf stage (growth stage 35) of the maize plants.</li> <li>Grey leaf spot - commence with treatment before 3 % of the total leaf surface area exhibit disease symptoms, or if symptoms are observed on the bottom 3 to 5 leaves.</li> <li>A follow-up spray might not be required in the case of <u>short season hybrids</u>, when the first spray was applied after pollination, as it may be sufficient to control the disease.</li> <li>Apply a maximum of two (2) <b>Mycoblock 250 SC</b> applications in the same season.</li> <li>If multiple applications are required to control the disease, the season's last application should be made using a registered fungicide from another chemical group, other than strobilurin (e.g. <b>SCOPE 250 EC</b>).</li> </ul> |  |
|   | <p>400 ml / ha<br/><b>Mycoblock 250 SC</b></p> <p><b>FOLLOWED BY</b></p> <p>300 ml / ha <b>SCOPE 250 EC</b><br/><b>PLUS</b><br/>a suitable oil adjuvant</p>  | <p>If the first treatment was applied within 21 days before pollination, or in the case of <u>medium to long-season hybrids</u>, apply a follow-up treatment with <b>SCOPE 250 EC</b> 3 to 5 weeks later.</p>  |
|   | <p>2 applications of<br/><b>Mycoblock 250 SC</b> at 300 ml / ha</p> <p><b>FOLLOWED BY</b></p> <p>300 ml / ha<br/><b>SCOPE 250 EC</b><br/><b>PLUS</b><br/>a suitable oil adjuvant</p>   | <p>Sometimes a third application may be required in cases where the first application was made very early. In such a case use <b>Mycoblock 250 SC</b> for the first two applications at 4 to 5 week intervals. Apply the last application 3 to 5 weeks after the second application using <b>SCOPE 250 EC</b>.</p> |
|   | <p><b>Row Application:</b></p> <ul style="list-style-type: none"> <li>In situations where maize is planted at wider row spacing's than 1 meter, e.g. in the Free State and western parts of RSA, row application may be used.</li> <li>For row applications, adjust dosage rates and spray volumes accordingly.</li> <li>The sprayer must be equipped with a tandem fitting of two hollow cone nozzles, mounted on the boom, directed on to each row.</li> <li>Apply in a sufficient volume of water to ensure good coverage of the plants.</li> </ul>   |  |



| <b>Crop &amp; Disease</b>   | <b>Dosage rate</b>  | <b>Remarks</b>  |
|---|---|---|
| <b>Potatoes</b><br>Early blight<br>( <i>A. solani</i> )   | 175 ml / ha<br><b>OR</b><br>250 ml / ha<br><b>OR</b><br>350 ml / ha | <b>Ground and Aerial application:</b><br>Add a Villa approved oil adjuvant to <b>SCOPE 250 EC</b> . Apply in a protective programme. Commence applications immediately before flowering (or earlier if the disease starts to develop) and continue until the plants start to senescence (end of flower drop). Apply the 175 ml per hectare dosage weekly, or the 250 ml per hectare dosage every 10 days, or the 350 ml per hectare dosage every 2 weeks. |
| <b>Soybeans</b><br>Soybean rust<br>( <i>P. pachyrhizi</i> )   | 325 ml / ha   | Apply at the first signs of disease and repeat 2 to 3 weeks later. If weather conditions favour disease development (temperatures of 15 to 28 °C and high humidity) use the shorter interval. If the disease appears during flowering or early pod-fill, more than one spray may be necessary. Should disease appear late in the growing season, a single application will be sufficient. Ensure thorough coverage of the whole plant.                    |
| <b>Table and Wine Grapes</b><br>Powdery mildew<br>( <i>O. tuckerii</i> )  | 125 ml / ha   | Apply every 14 days in a spray programme as a full cover spray to the point of run-off. Commence spraying when the first shoots are 100 to 250 mm long and repeat the treatment at intervals as indicated.<br><b>Do not apply more than 4 treatments per season. The last application must not be later than bunch closure (berry touch completed).</b><br>Always ensure thorough coverage.   |
| <b>NOTE</b>   |   |   |
| <ul style="list-style-type: none"> <li>For <b>wine grapes</b> increase the spray volume progressively, from 250 litres per hectare to reach 1000 litres per hectare at pea-berry stage and repeat at this rate throughout the rest of the season.</li> <li>For <b>table grapes</b> the spray volume should be increased progressively from 500 litres per hectare to 1200 litres per hectare</li> <li><b>Do not exceed the spray interval.</b></li> </ul>   |   |   |
| <b>Tomatoes</b><br>Early blight<br>( <i>A. solani</i> )   | 35 ml / 100 l water   | <b>High volume:</b> Apply at 500 to 2000 litres spray mixture per hectare.  |
|   | 350 ml / ha   | <b>Low volume:</b> Mistblower application.  |
| <b>NOTE</b>   |   |   |
| <ul style="list-style-type: none"> <li>For <b>low Early blight pressure areas where other diseases are not a problem:</b> <ul style="list-style-type: none"> <li>Apply <b>SCOPE 250 EC</b> at 14-day intervals.</li> </ul> </li> <li>For <b>moderate to heavy Early blight pressure areas and/or where other diseases need to be controlled:</b> <ul style="list-style-type: none"> <li>Alternate weekly between <b>SCOPE 250 EC</b> and a suitable broad-spectrum fungicide (e.g., <b>Mancozeb</b>, <b>Chlorothalonil</b> or <b>Copper</b>).</li> <li>Commence applications from the flowering stages (about 8 weeks after transplant) or earlier if the disease already started developing. Continue with <b>SCOPE 250 EC</b> into the picking stages (16 to 20 weeks after transplant) applying a maximum of eight (8) <b>SCOPE 250 EC</b> applications per season.</li> <li>During the pre-flowering stages and after the <b>SCOPE 250 EC</b> treatments are discontinued, apply a suitable broad-spectrum fungicide.</li> </ul> </li> <li>Ensure complete coverage of the entire plant mass with spray mixture at all times.</li> <li><b>SCOPE 250 EC</b> does not control Late blight (<i>Phytophthora infestans</i>).</li> </ul> |   |   |

The following products mentioned in this label can be replaced with the following equivalent products:

- MYCOBLOCK 250 SC** (L 8591) = **STROBIN 250 SC** (L 8625) (**Azoxystrobin**),
- INSPECT 400 SC** (L 9182) = **DETECT 400 SC** (L 9238) = **EXPOSE 400 SC** (L 9511) (**Dithianon**)  
and
- VILLA UNIZEB 800 WP** (L 8056 / N-AR 1091) = **COZEB 800 WP** (L 7098) = **NEWZEB 800 WP** (L 8167) (**Mancozeb**).

**MYCOBLOCK 250 SC, COZEB 800 WP and/en DETECT 400 SC** are registered products of / is  
geregistreeerde produkte van  
**Universal Crop Protection (Pty) Ltd.**

**INSPECT 400 SC and/en VILLA UNIZEB 800 WP** are registered products of / is geregistreeerde  
produkte van  
**Villa Crop Protection (Pty) Ltd.**

**NEWZEB 800 WP, STROBIN 250 SC and/en EXPOSE 400 SC** are registered products of / is geregistreeerde  
produkte van  
**Cropasure (Pty) Ltd.**