

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Insecticide



ACEPHATE 750 SP

Reg. No. L 7181 Act/Wet No. 36 of/van 1947
N-AR 1929

2: 05/05/09-Jan2019

A systemic and contact water-soluble insecticide for the control of pests mentioned on the crops listed.

'n Sistemiese en kontak wateroplosbare insekdoder vir die beheer van die plae vermeld op gewasse soos aangedui.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL
acephate (organophosphate) **750 g/kg** asefaat (organofosfaat)

IRAC INSECTICIDE GROUP CODE **1B** IRAC INSEKDODER GROEPKODE

kg

villa

Registration holder / Registrasiehouer:

Villa Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1992/002474/07

PO Box / Posbus 10413, Aston Manor, 1630

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: 2783

Willow Set & Print 011 394-4486



CAUTION
VERSIGTIG



GEBRUIKSAANWYSINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED
REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

ACEPHATE 750 SP

Reg. No. L7181 / N-AR 1929 Act/Wet No. 36 of/van 1947

IRAC INSECTICIDE GROUP CODE / IRAC INSEKODER GROEPKODE: 1B

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

acephate (organophosphate) / asefaat (organofosfaat) 750 g/kg

Registration holder / Registrasiehouer:

VILLA CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1992/002474/07 Mpy. Reg. Nr.

PO Box / Posbus 10413,

ASTON MANOR, 1630, Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS****Withholding periods:**

Minimum time between the last application and harvest:	
Apples and Pears	30 days
Cruciferae and Tomatoes	3 days
Potatoes	14 days
Peaches and Plums	28 days
Table and Wine Grapes	40 days
Tobacco	14 days

IN THE CASE OF CROPS INTENDED FOR EXPORT, CONFIRM WITHHOLDING PERIODS WITH THE RELEVANT EXPORT ORGANISATION. IF THIS PRODUCT IS USED ACCORDING TO THE RECOMMENDATIONS ON THIS LABEL, LOCAL MRL VALUES WILL NOT BE EXCEEDED. HOWEVER, VILLA CROP PROTECTION (PTY) LTD. CANNOT ACCEPT RESPONSIBILITY FOR EXPORTED CROPS EXCEEDING THE IMPORT TOLERANCES OF OTHER COUNTRIES.

- Handle with care.
- May irritate skin and eyes.
- Toxic to fish, bees and wild life.
- Store in a cool, dry place away from food, feeds, fertilizers and other agricultural remedies.
- Keep out of reach of children, animals and uninformed persons.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Aerial application:

Notify all inhabitants of the immediate area to be sprayed and issue the necessary warnings. Do not spray over, or allow drift to contaminate water of adjacent areas.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Do not inhale the powder dust and spray mist.
- Avoid contact with eyes and prolonged or repeated skin contact.
- Wash contaminated clothing after use.
- Wash with soap and water after use or accidental skin contact.
- Do not eat, drink or smoke whilst mixing, applying the product or before washing hands and face or change of clothing.
- Avoid drift of spray onto other crops, grazing, rivers, dams and areas not under treatment.
- Clean applicator after use and before using it for other remedies.
- Dispose of wash water where it will not contaminate crops, grazing, rivers and dams.

- Prevent contamination of food, feed, drinking water and eating utensils.
- Invert the empty container over the spray or mixing tank to make sure the container is empty before destroying the container.
- Destroy the empty container by perforation and flattening.
- **Never** re-use the container for any other purpose.

RESISTANCE WARNING

ACEPHATE 750 SP is a group code 1B insecticide. Any insect population may contain individuals naturally resistant to **ACEPHATE 750 SP** and other group code 1B insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **ACEPHATE 750 SP** or any other group code 1B insecticide.

To delay insecticide resistance:

- avoid exclusive repeated use of insecticides from the same insecticide group code. Alternate or tank mix with products from different insecticide group codes,
- integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

ACEPHATE 750 SP should be tested on a small scale when mixed with other pesticides, to ensure that it is safe to the crop.

Mixing instructions:

- Half fill the spray tank with water and add the required amount of **ACEPHATE 750 SP** to the water, while agitating.
- Ensure that all the lumps have dissolved before adding the rest of the water.
- Maintain agitation during mixing and application.

General directions:

Use at all times correctly calibrated and suitable equipment in good working condition, to ensure wetting of all parts of the plant thoroughly.

Aerial application:

Aerial application of **ACEPHATE 750 SP** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- Volume: A spray mixture volume of 30 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.
- Droplet coverage: 25 to 35 droplets per cm² must be recovered at the target area.
- Droplet size: A droplet spectrum with a VMD of 280 to 300 micron is recommended. Limit the production of fine droplets less than 150 micron (high drift and evaporation potential) to a minimum.
- Flying height: Maintain the height of the spray boom at 3 to 4 metres above the target. Do not spray when aircraft dives, is in a climb or when banking.
- Use suitable atomizing equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomizers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8°C.
- Stop spraying if the wind speed exceeds 15 km per hour.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),

- b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

APPLICATION RATES

Crop / Pest	Dosage Rate	Remarks
Apples and Pears Codling moth	50 g / 100 ℓ water	Apply as a full cover spray at 75 % petal fall and repeat 14 days later. Do not apply more than two sprays per season. The last spray must be applied before the end of November.
African bollworm	50 g / 100 ℓ water	Apply as a full cover spray and repeat if necessary. Do not apply more than two sprays per season. The last spray must be applied before the end of November.
Banded fruit weevil (Snout beetle)	50 g / 100 ℓ water	Apply as a full cover spray between 3/4 petal drop and at the end of November. Sprays should be repeated after 14 days. Do not apply more than two sprays per season.
Leafroller, Antestia & Fruit nibbler	50 g / 100 ℓ water	Application sprayed for the control of Codling moth, Snout beetle and African bollworm will also control Leafroller, Antestia and Fruit nibbler present at application. Do not apply more than two sprays. The last spray must be applied before the end of November.
Cruciferae Diamond back moth larvae	350 to 500 g / ha	Apply as a full cover spray when the pest is noticed. Ensure complete coverage of plants, especially the underside of leaves. The addition of a wetting agent is recommended. Repeat every 7 to 10 days. The lower rate should be repeated every 7 days and the higher rate is recommended for the first application.
Plums African bollworm, Antestia bug, Fruit nibbler, Banded fruit weevil (Snout beetle), Leafroller & Codling moth	50 g / 100 ℓ water	Western Cape only: Apply as a full cover spray, to ensure thorough wetting and repeat as necessary. Do not apply more than two sprays per season. Do not apply during blossom, as it is toxic to bees. Do not spray after petal fall.
Peaches Dessert and Canning Aphids	25 g / 100 ℓ water <u>High Volume:</u> 0.38 to 0.63 kg / ha	Western Cape only: Apply as a full cover spray when the pest is noticed and repeat as necessary. Do not apply more than two sprays per season. If applications coincide with the Oriental fruit moth day degree model, Oriental fruit moth may be suppressed but not controlled.
Antestia bug, Codling moth, Fruit nibbler & Banded fruit weevil (Snout beetle)	50 g / 100 ℓ water <u>High Volume:</u> 1.25 to 1.75 kg / ha <u>Low Volume:</u> 1.0 to 1.4 kg / ha	Western Cape only: Apply as a full cover spray when pest is noticed and repeat as necessary. Do not apply more than two sprays per season. Do not apply during blossom, as it is toxic to bees. Do not spray after petal fall. Will also control African bollworm and Fruit nibbler.
African bollworm	50 g / 100 ℓ water <u>High Volume:</u> 1.25 to 1.75 kg / ha <u>Low Volume:</u> 1.0 to 1.4 kg / ha	Western Cape only: Apply at 75% petal fall and repeat 14 days later. Do not apply more than two sprays per season. Do not apply during blossom, as it is toxic to bees. Will also control Codling moth, Antestia bug, Fruit nibbler and Weevil.

Crop / Pest	Dosage Rate	Remarks
Table and Wine Grapes Weevil (Snout beetle)	50 to 75 g / 100 ℓ water <u>High Volume:</u> 0.13 to 1.1 kg / ha	Apply as a full cover spray at 250 to 1500 litres spray mixture per hectare, as soon as the pest is noticed. Use higher dosage rate in cases of severe infestation. Repeat 7 to 14 days later if necessary.
Potatoes Potato tuber moth	500 g / ha	Apply as a full cover spray in 200 to 500 litres water per hectare, depending on plant size. Start application when plants are one month old and repeat every 7 to 14 days if necessary. This programme will also control Aphids. Ridge at least twice during the season.
Aphids	500 to 750 g / ha	Apply at first sign of infestation and repeat every 7 days. When the Aphid infestation is high, use the higher rate to ensure initial control. Apply as a full cover spray in 200 to 500 litres water per hectare, depending on plant size. Aerial application: Apply in 30 litres of water per hectare.
Tobacco African bollworm & Leaf miner	200 g / 100 ℓ water	Ground application: Apply as a full cover spray. Where a heavy infestation occurs, a second application is recommended 7 days later. Repeat if necessary. Ensure thorough coverage of all leaves.
African bollworm	1000 g / ha	Aerial application (also follow the instructions under "DIRECTIONS FOR USE"): 1. Apply only when larvae are smaller than 12.5 mm. 2. Repeat application after 7 days, if necessary repeat application again. 3. Application must preferably be performed in the morning before the larvae migrate to the lower leaves. 4. Application to high-density tobacco is not recommended.
Slug	200 g / 100 ℓ water	Apply as a full cover spray and repeat application every 7 days up to 14 days before harvest. When a heavy infestation occurs, apply 400 g ACEPHATE 750 SP per 100 litres water as a full cover spray and re-apply 200 g ACEPHATE 750 SP per 100 litres water, 7 days later.
Cutworm	165 g / 100 ℓ water	Ground application: Pour 30 ml of the solution over each young plant. Repeat if necessary.
Green aphid (<i>Myzus persicae</i>)	200 g / 100 ℓ water	Apply as a full cover spray and ensure thorough coverage of all leaves. Where a heavy infestation occurs, a second application is recommended 7 days after the first application, to obtain good initial control. Repeat application if necessary every 7 days.
Tobacco Seed beds Cutworm & Leaf eating insects	1 g / m ²	Ground application: 1) Spray: Mix 200 g ACEPHATE 750 SP per 100 litres water and use a coarse spray at the rate of 500 ml solution per m ² every 7 days, to get good run-off onto the soil. OR 2) Drench: Mix 65 g ACEPHATE 750 SP per 100 litres water and use a watering can to apply 1.5 litres solution per m ² , every 7 days.

Crop / Pest	Dosage Rate	Remarks
Tomatoes African bollworm	75 to 100 g / 100 ℓ water	Apply a minimum of 1000 litres spray mixture per hectare and increase this spray volume to a maximum of 2000 litres per hectare, according to plant size. Ensure thorough coverage of plants. Commence spraying at flowering and repeat application every 7 days, as long as pest remains active. Use the higher rate as a first treatment, when infestation is present. Repeat application after rain.
Thrips	100 g / 100 ℓ water	Apply as a full cover spray and ensure thorough coverage of plants. Where a heavy infestation occurs, a second application is recommended 7 to 10 days after first application to obtain good initial control. If necessary, repeat application every 7 to 10 days.