

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Insecticide



SEVIN® XLR PLUS

Reg. No. L 5783 Act/Wet No. 36 of/van 1947

2:12/10/2015-Jan2017

*A residual contact and stomach insecticide formulated as a suspension concentrate, for the control of various insects on crops, in gardens and lawns, hides and skins as indicated and for fruit thinning in apples.*

*'n Nabywende kontak- en maaggif geformuleer as 'n suspensie konsentraat, vir die beheer van verskeie plae op gewasse, in tuine en grasperke, huide en velle asook vir vruguitdunning in appels.*

**ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL**

carbaryl (carbamate) **480 g/l** karbariel (karbamaat)

IRAC INSECTICIDE GROUP CODE **1A** IRAC INSEKDODER GROEPKODE



**villa**

Registration holder / Registrasiehouer:  
Universal Crop Protection (Pty) Ltd.  
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07  
PO Box / Posbus 801, Kempton Park, 1620  
Tel: 011 396 2233  
Website / Webblad: [www.villacrop.co.za](http://www.villacrop.co.za)

UN Number: 3082



**HARMFUL  
SKADELIK**

Willow Set & Print 011 394-4486



GERUUKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE  
GEDRUK OP HOUER/SAK

Date formulated:

Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED  
ON CONTAINER/BAG

Batch number:

Lotnummer:

**SEVIN® XLR PLUS**

Reg. No. L 5783 Act/Wet No. 36 of/van 1947

IRAC INSECTICIDE GROUP CODE / IRAC INSEKODER GROEPKODE: 1A

**ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:**

carbaryl (carbamate) / karbariel (karbamaat).....480 g/l

Registration holder / Registrasiehouer:

**UNIVERSAL CROP PROTECTION (PTY) LTD.**

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801

KEMPTON PARK, 1620

Tel. (011) 396 2233

**HARMFUL****SKADELIK****WARNINGS****Withholding periods:**

<b>Allow the following periods between the last application and harvest and/or feeding of the crops concerned:</b>	
Apples & Pears	7 days
Cotton (Grazing)	7 days
Grapes (Table) Do not apply after berries have reached pea size (4 to 6 weeks prior to harvest), as this may cause tarnishing.	28 to 42 days
Grapes (Wine)	14 days
Litchi, macadamia & Pecan nut trees (Stem application)	0 days
Lucerne (Alfalfa) (Grazing)	14 days
Maize	14 days
Pastures & Field crops (Harvesting and grazing)	14 days
Prickly pear & Spineless cactus ( <i>Opuntia</i> spp.)	1 day
Tobacco	14 days

**NOTE**

**IF THE CROP TO BE TREATED IS INTENDED FOR EXPORT, CONSULT THE RELEVANT IMPORTER OR EXPORTING BODY REGARDING THE USE OF THIS PRODUCT, MAXIMUM RESIDUE LIMITS AND RECOMMENDED WITHHOLDING PERIODS.**

- **DO NOT APPLY TO KIEFFER OF FORELLE PEAR VARIETIES OR QUINCES.**
- **DO NOT APPLY TO GRANNY SMITH OR VIGOROUS ROYAL GALA TYPE APPLE TREES IF CONDITIONS FAVOUR FRUIT THINNING.**
- **DO NOT APPLY TO GOLDEN DELICIOUS APPLE TREES IN AREAS WHERE RUSSETING IS A PROBLEM.**
- Handle with care.
- Poisonous when swallowed or inhaled.
- Toxic to fish, bees and wildlife.
- Store in a cool, dry place away from food and feeds.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated field within 1 day after application unless wearing protective clothing.
- **In case of poisoning immediately call a physician and make this label available to him/her.**

**Aerial Application:**

Notify all inhabitants of the immediate area to be sprayed and issue the necessary warnings. Avoid overlapping of applications and do not spray over or allow drift to contaminate water or adjacent areas.

**ALTHOUGH SEVIN® XLR PLUS IS REGARDED AS CROP-SAFE ON MOST IMPORTANT CULTIVARS; THIS DOES NOT MEAN THAT A MORE SENSITIVE CULTIVAR MIGHT NOT BE COMMERCIALIZED IN THE FUTURE. WHERE A NEW CULTIVAR IS ENCOUNTERED, LARGE AREAS SHOULD NOT BE SPRAYED WITHOUT PRIOR TESTING OF SEVIN® XLR PLUS ON THAT CULTIVAR.**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions. The action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

### **PRECAUTIONS**

- Avoid prolonged or repeated skin contact with concentrate or spray.
- Do not inhale fumes or the spray mist.
- Wear a face shield and rubber gloves while handling and applying the product.
- Wash contaminated clothing daily.
- Wash with soap and water immediately after use or accidental skin contact.
- Prevent contamination of food, eating utensils, feeds and drinking water.
- In case of eye splashes, rinse thoroughly with clean water.
- Do not eat, drink or smoke whilst mixing or applying, or before washing hands and face and change of clothing.
- Do not spray over or allow drift to contaminate water bodies such as dams, ponds, rivers, streams or fish hatcheries.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of one third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the container by perforation and flattening and dispose of it in a safe way.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

### **SYMPTOMS OF POISONING**

Symptoms may include salivation, watery eyes, pinpoint eye pupils, blurred vision, muscle weakness, difficult breathing, excessive sweating, abdominal cramps, nausea, vomiting, diarrhoea, weakness and headaches. In severe cases, convulsions, unconsciousness and respiratory failure may occur.

### **FIRST AID TREATMENT**

- Remove patient from source of poisoning and keep him/her quiet and reassured.
- **Skin:** Remove contaminated clothing, shoes and leather goods. Wash skin gently and thoroughly with cold water and non-abrasive soap. Do not rub skin. Obtain medical attention if irritation persists.
- **Eyes:** Flush eyes with clean water for at least 15 to 20 minutes, holding the eyelid(s) open. Remove contact lenses, if present, after the first 5 minutes then continue rinsing eye(s). Obtain medical attention if irritation persists.
- **Inhalation:** Maintain airway and administer oxygen if available. Remove the patient from the source of contamination and keep affected person warm and at rest. Get medical attention immediately.
- **Ingestion:** Obtain medical attention immediately and make this label available to him or her. Do not give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by tickling the back part of the throat with your finger. Repeat until vomit is clear and free from smell of poison. Keep under medical supervision.

### **NOTE TO PHYSICIAN**

**SEVIN® XLR PLUS (Carbaryl)** is an oxime carbamate which is a moderate reversible, cholinesterase inhibitor. Gastric lavage may be used if this product has been swallowed. Specific treatment consists of atropine sulphate. Improve tissue oxygenation as much as possible before administering atropine to minimise risk of ventricular fibrillation. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinisation has been achieved and repeated thereafter whenever symptoms re-appear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly

every 30 to 60 minutes as needed to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from over-exposure is to be expected within 24 hours. Narcotics and other sedatives should not be used. Other drugs like 2-PAM are not recommended.

### **RESISTANCE WARNING**

**SEVIN® XLR PLUS** is a group code 1A insecticide. Any insect population may contain individuals naturally resistant to **SEVIN® XLR PLUS** and other group code 1A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **SEVIN® XLR PLUS** or any other group code 1A insecticide.

Since the presence of resistant insects is difficult to detect prior to insecticide application, it is of the utmost importance that treated areas be inspected at regular intervals to identify the occurrence of insecticide resistant insects timeously. The registration holder will not accept liability for performance failures in the event of a build-up of resistant insects resulting from inadequate resistance management practices as indicated on the label.

To delay insecticide resistance:

- avoid exclusive repeated use of insecticides from the same insecticide group code. Alternate or tank mix with products from different insecticide group codes,
- integrate other control methods (chemical, cultural, biological) into insect control programmes.

### **Codling moth resistance management**

**When used to control Codling moth (*Cydia pomonella*) in pome fruit, a maximum of four (4) SEVIN® XLR PLUS sprays must be applied to a single Codling moth generation.**

**To prevent or delay the development of resistance, products with other modes of action should be applied for the control of the other two generations. If SEVIN® XLR PLUS is used for the control of the last generation of the season a product with another mode of action should be used for the control of the first generation in the next season.**

For specific information on **RESISTANCE MANAGEMENT** contact the registration holder of this product. For the period that has to be allowed between last application and harvest or grazing, refer “**WARNINGS**”.

### **DIRECTIONS FOR USE: Use only as directed.**

#### **Compatibility:**

**SEVIN® XLR PLUS** is compatible with **PLANOFIX®** (Reg. L 531). Do not mix with other chemicals when used for fruit thinning. Do not mix with highly alkaline materials e.g. **Hydrated lime** or **Lime sulphur**.

#### **Mixing instructions:**

- Half fill the spray tank with clean water.
- Add the required quantity of **SEVIN® XLR PLUS** as recommended below and top up to the volume of water required. Ensure thorough agitation.
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

#### **Aerial application:**

**SEVIN® XLR PLUS** can be applied aurally on certain crops where this is indicated in the “**APPLICATION RATES**” below.

**DO NOT COMMENCE AERIAL APPLICATION BEFORE CONSULTING YOUR SEVIN® XLR PLUS REPRESENTATIVE.**

Aerial application of **SEVIN® XLR PLUS** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- **Volume:** A minimum spray mixture volume of 30 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aurally at a lower volume rate than recommended above.
- **Droplet coverage:** 30 to 40 droplets per cm<sup>2</sup> must be recovered at the target area.
- **Droplet size:** A droplet spectrum with a VMD of 250 to 280 micron is recommended. Limit the production of fine droplets, less than 150 micron (high drift and evaporation potential), to a minimum.

- **Flying height:** Maintain the height of the spray boom at 3 to 4 metres above the target. Do not spray when aircraft dives, is in a climb or when banking.
- Use suitable **atomizing equipment** that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomizers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the **wingtip vortices**.
- The difference in **temperature** between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8°C.
- Stop spraying if the **wind speed** exceeds 15 km per hour.
- Stop spraying under **turbulent**, unstable and dry conditions during the heat of the day.
- Spraying under temperature **inversion conditions** (spraying in or above the inversion layer) and/or **high humidity conditions** (relative humidity 80 % and above) may lead to the following:
  - a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),
  - b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

#### APPLICATION RATES

<b>Crop / Pest</b>	<b>Dosage rate</b>	<b>Remarks</b>
<b>Apples</b> <b>Chemical thinning of fruit</b>  <u>Easy-to-Thin cultivars</u> e.g., Granny Smith, Standard red cultivars and Young, vigorous Royal Gala types	Use on Granny Smith and young Royal Gala types only under expert guidance and only if conditions do not favour fruit thinning, otherwise over thinning might result. On all of these cultivars <b>SEVIN® XLR PLUS</b> must not be combined with, or sprayed in a programme with other thinning remedies. <b>Refer also to "NOTES"</b> .	
	100 to 150 ml / 100 ℓ water	For early thinning, apply once as a full cover spray in the period between 80 and 100 % petal fall.
	25 to 100 ml / 100 ℓ water	For later thinning, apply once as a full cover spray in the period between 10 and 15 mm fruit diameter.
<u>Difficult-to-Thin cultivars</u> e.g., Red spur types, Braeburn, Cripps' Red, Cripps' Pink, full-bearing Royal Gala types, Golden Delicious types and Fuji types	Use on Golden Delicious only in areas where fruit russetting is not a problem. <b>Refer also to "NOTES"</b> .	
	100 to 150 ml / 100 ℓ water	For early thinning under conditions that favour fruit thinning, apply once as a full cover spray in the period between 80 and 100 % petal fall.
	100 to 150 ml <b>SEVIN® XLR PLUS</b>  <b>PLUS</b>  11 ml <b>Planofix®</b> / 100 ℓ water	Under conditions that do not favour fruit thinning, apply a tank mixture of <b>SEVIN® XLR PLUS</b> and <b>Planofix®</b> once as a full cover spray in the period between petal fall and 12 mm fruit diameter.
	100 to 150 ml / 100 ℓ water	As an alternative to the above treatment, if conditions do not favour fruit thinning, a <b>SEVIN® XLR PLUS</b> treatment can be preceded by a <b>Planofix®</b> application between petal fall and 5 mm fruit diameter. Follow this with an application of <b>SEVIN® XLR PLUS</b> at 10 to 15 mm fruit diameter.

<b>Crop / Pest</b>	<b>Dosage rate</b>	<b>Remarks</b>
<b>Apples &amp; Pears</b> (Refer “ <b>WARNINGS</b> ” for varieties that must not be treated)	<b>IMPORTANT</b> Refer “ <b>Warning against Codling moth resistance</b> ” under “ <b>RESISTANCE WARNING</b> ”.	Commence application at the beginning of a moth generation or at 75 % petal fall (in the case of the first generation). Repeat at 14 to 18 day intervals for a maximum of four applications.
Codling moth	100 mℓ / 100 ℓ water	Apply as a full cover spray at 2 weekly intervals. To avoid possible apple thinning use an alternative registered Codling moth remedy for the first application, where after <b>SEVIN® XLR PLUS</b> can be used. Do not use <b>SEVIN® XLR PLUS</b> within 21 days after full bloom.
Leaf rollers	225 mℓ / 100 ℓ water	When used in Codling moth programme, <b>SEVIN® XLR PLUS</b> will control Leaf rollers. However, where a single treatment is required, the dosage must be increased as prescribed. Apply as a full cover spray when pest is noticed.
Pear bud mite	100 mℓ / 100 ℓ water	When used in Codling moth programme, <b>SEVIN® XLR PLUS</b> will control Pear bud mite provided an additional treatment is carried out by the 3 <sup>rd</sup> week of November as a full cover spray.
Mealybugs	225 mℓ / 100 ℓ water	Apply as a full cover spray. Commence treatment early December and follow-up with 1 to 2 additional treatments depending on the degree of infestation. These treatments will coincide with the normal Codling moth programme. <b>Do not</b> apply later than mid-January, as later applications will not control this pest satisfactorily.
<b>Castor Oil</b> Castor oil looper ( <i>Achaea</i> spp.)	225 mℓ / 100 ℓ water	<u>LV</u> : Commence spraying when infestation is first noticed.
<b>Cotton</b> Red & Spiny bollworm, Stainers, Elegant grasshoppers, Syagrus beetles, Aphids, Leafhoppers, Leaf-eaters ( <i>Anomis</i> , <i>Xanthodes</i> )	Commence cotton spraying when first buds or “squares” appear and continue treatments at 7 to 10 day intervals until the first bolls open. Re-spray if 13 mm or more rain falls within 24 hours of treatment.	
	<u>Tail boom</u> : 1050 mℓ / 100 ℓ water	For tail boom apply 50 to 250 litres of the spray mixture per hectare depending on plant size.
	<u>Aerial application</u> : 1325 to 2650 mℓ / ha	Apply 1325 mℓ per hectare to cotton up to 60 cm in height or approximately 50 to 60 % canopy and thereafter 2650 mℓ per hectare in not less than 30 litres water per hectare. (Refer detailed “ <b>Instructions</b> ” above).
<b>Field Crops</b> Elegant grasshoppers	225 mℓ / 100 ℓ water <b>OR</b> 2.0 mℓ / 10 m <sup>2</sup>	Apply preferably to young stages.
Army worm	550 mℓ / ha <b>OR</b> 2.0 mℓ / 10 m <sup>2</sup>	<u>LV</u> : Apply as a cover spray and commence treatment as soon as infestation is noticed. <u>Aerial application</u> : Apply in not less than 30 litres water per hectare. <b>Refer to “Aerial Application”.</b>

<b>Crop / Pest</b>	<b>Dosage rate</b>	<b>Remarks</b>
<b>Grapes</b> (Table and wine) Weevils (Snout-beetles)	225 ml / 100 l water <b>OR</b> 23 ml / 10 l water	When applied against Mealybugs, <b>SEVIN® XLR PLUS</b> will assist in the control of vine Weevils.
Mealybugs	225 ml / 100 l water <b>OR</b> 23 ml / 10 l water	Commence spraying as soon as the pest is noticed at the bases of young shoots. Repeat application every 3 weeks if necessary.
<b>Gardens &amp; Lawns</b> Lawn caterpillar	800 ml / 100 l water <b>OR</b> 2.0 ml / 10 m <sup>2</sup>	Apply as a full cover application.
<b>Litchi, Macadamia nut &amp; Pecan nut trees</b> Stem and bark borers	450 ml / 100 l water	Apply approximately 50 ml of spray mix directly into each tunnel by means of a knapsack applicator for control of Stem borer or spray the lesions for bark borer control. Do not spray the entire tree. Treat only the affected areas on the stem.
<b>Lucerne</b> Lucerne caterpillar	1100 to 1500 ml / ha	Dosage depends on plant density and plant size. <u>LV</u> : Full cover application. <u>Aerial application</u> : Apply in not less than 30 litres water per hectare. <b>Refer to “Aerial Application”.</b>
<b>Maize</b> Maize chafer beetles	225 ml / 100 l water <b>OR</b> 23 ml / 10 l water	Apply as a full cover spray and repeat when necessary.
<b>Prickly Pear &amp; Spineless Cactus</b> ( <i>Opuntia</i> spp.)  <i>Cactoblastis cactorum</i>	300 ml / 100 l water <b>OR</b> 30 ml / 10 l water	Apply as a coarse spray by means of a knapsack sprayer fitted with D5 nozzles. Add a wetting agent. Spray the entire plant surface to the point of run off, especially egg sticks. Commence treatment during egg stage (two generations per annum) i.e. for the Karoo area, normally during the first two weeks of March and November. For other areas treatment should commence after oviposition (egg-laying) peak, but before hatching of eggs.
<i>Cochineal</i> ( <i>Dactylopius</i> spp.)	300 ml / 100 l water <b>OR</b> 30 ml / 10 l water	Apply as for <i>Cactoblastis</i> . Commence treatment when infestation is noticed and repeat if necessary.
<b>Tobacco</b> American bollworm, Tobacco slug & Army worm	<u>Ground application</u> : 225 ml / 100 l water  <u>Aerial application</u> : 1750 ml / ha	Do not use against Tobacco slug if tobacco plants are fully grown.  <u>HV</u> : Apply as a full cover application as soon as pest is noticed.  <u>Aerial application</u> : Apply as soon as pest is noticed in not less than 30 litres water per hectare. <b>Refer to “Aerial Application”.</b>
<b>Hides &amp; Skins</b> Domesticated hide and skin beetle larvae	450 ml / 100 l water <b>OR</b> 45 ml / 10 l water	Apply generously as a coarse spray or by means of a paint brush to hides and skins as well as to sheltering places such as sacking and cracks and crevices in floors and walls. Pay special attention to the flesh side of hides. Five (5) litres will be sufficient to treat approximately 50 skins or 15 hides.

**NOTES**

- Expert advice should be sought when using **SEVIN® XLR PLUS** for the first time to thin apples.

- Apple tree reaction to chemical fruit thinners is dependent on many interacting variables such as the degree of thinning required, environmental conditions, cultivar, rootstock, tree age, tree shape and the use of other thinning agents. Local knowledge of the abovementioned factors is essential if the desired result is to be attained.
- **SEVIN® XLR PLUS** must **NOT** be used on Golden Delicious strains in areas where retiform russetting is a problem.
- On Royal Gala up to 5 years old, apply only a single fruit thinning treatment of **Planofix®** or **SEVIN® XLR PLUS**.

**PLANOFIX®** is a registered trade mark of / is 'n geregistreerde handelsmerk van  
**BAYER (PTY) LTD.**

**SEVIN® XLR PLUS** is a registered trade mark of / is 'n geregistreerde handelsmerk van  
**TESSENDERLO KERLEY INC.**