

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Micronutrient



MAX-IN[®] SULFUR

Reg. No. B 5334 Act/Wet No. 36 of/van 1947

1: 19/09/2017 – June 2018

For use in a balanced fertilizer program for all crops requiring sulfur and potassium.

Vir gebruik in 'n gebalanseerde misstof program op alle gewasse wat swael en kalium benodig.

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE

sulfur / swael (S) 130 g/kg
potassium / kalium (K) 163 g/kg

SG: 1.34 at 20 °C

FERTILIZER GROUP/GROEP 2 MISSTOF



villa

Registration holder / Registrasiehouer:
Winfield Solutions Registration Holdings (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 2015/312008/07
PO Box / Posbus 10413, Aston Manor, 1630
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: Not regulated

GEBRUIKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

MAX-IN® SULFUR

Reg. No. B 5334 Act No. 36 of 1947

FERTILIZER GROUP/GROEP 2 MISSTOF

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE:

sulfur / swael (S)	130 g/kg
potassium / kalium (K)	163 g/kg

Registration holder / Registrasiehouer:

WINFIELD SOLUTIONS REGISTRATION HOLDINGS (PTY) LTD.

Co. Reg. No. 2015/312008/07 Mpy. Reg. Nr.

P.O. Box / Posbus 10413, Aston Manor, 1630,

Tel. (011) 396 2233

DIRECTIONS FOR USE

- For use in a balanced fertilizer program for all crops requiring potassium and sulfur.
- User should seek the advice of a certified crop advisor or trained agronomist.
- For specific crop recommendations in your area, consult your state's recommendations.
- Use in a minimum of 100 litres of water per hectare.

CAUTION: Plant and leaf injury may occur on some crops when certain weather and growing conditions are present. Do not apply to foliage of any crop when temperatures are above 32°C.

MIXING INSTRUCTIONS

- Add water to the mixing tank and agitate.
- Add **MAX-IN® SULFUR**.

NOTE Care must be taken as this product foams during the mixing process. It is recommended to add **MAX-IN® SULFUR** just before final tank mixture volume is reached. Allow for complete dispersion before application.

- At the end of the spraying operation, thoroughly rinse out spray equipment and spray booms.

APPLICATION RATES

Crop	Application rate	Remarks
<u>Lucern</u>	2.25 ℓ to 4.5 ℓ/ha	Apply MAX-IN® SULFUR after each cutting. Applications should be made when sufficient regrowth has occurred.
<u>Cereal Crops</u>	2.25 ℓ to 4.5 ℓ/ha	Apply when plants are between tillering and early joint stages. Later application may be needed; tissue samples should be used to determine later usages.
<u>Maize</u>	2.25 ℓ to 4.5 ℓ/ha	Apply when corn is in the (V4 to V7) stage. Tissue samples should be used to determine usages.
<u>Other Crops</u>	-	Apply according to directions of your regional Villa Crop Protection Nutrition Specialist or local agricultural professional to crops where the addition of potassium and sulfur would be beneficial.

MAX-IN® SULFUR is a registered trademark of / is 'n geregistreerde handelsmerk van
WINFIELD SOLUTIONS REGISTRATION HOLDINGS (PTY) LTD.