

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Insecticide



UNIPEX 500 WDG

Reg. No. L 8474 Act/Wet No. 36 of/van 1947
W 130464

2:10/4/2019-April2019

A water dispersible contact granular insecticide for the control of aphids in crops as indicated.

'n Waterdispergeerbare kontak korrelinsekdoder vir die beheer van plantluise in gewasse soos aangedui.

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL
pirimicarb (carbamate) 500 g/kg pirimikarb (karbamaat)

IRAC INSECTICIDE GROUP CODE **1A** IRAC INSEKODER GROEPKODE

g

villa

Registration holder / Registrasiehouer:
Universal Crop Protection (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07
PO Box / Posbus 801, Kempton Park, 1620
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: 2757

Willow Set & Print 011 394-4486



HARMFUL
SKADELIK



GERUUKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE
Date formulated: GEDRUK OP HOUER/SAK
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED
ON CONTAINER/BAG
Batch number:
Lotnummer:

UNIPEX 500 WDG

Reg. No. L 8474 / W 130464 Act/Wet No. 36 of/van 1947
 IRAC INSECTICIDE GROUP CODE / IRAC INSEKODER GROEPKODE: 1A

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

pirimicarb (carbamate) / pirimikarb (karbamaat) 500 g/kg

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620

Tel. (011) 396 2233



HARMFUL SKADELIK

WARNINGS**Withholding periods:**

Minimum number of days between the last application and harvest:	
Grain sorghum	30 days
Peaches and Groundnuts (hay)	21 days
Pecan nuts	56 days
Potatoes, Citrus, Apples and Cotton	14 days
Wheat, Oats and Cruciferae	7 days

NOTE

COMPLIANCE WITH THESE WITHHOLDING PERIODS WILL ENSURE THAT RESIDUES DO NOT EXCEED LOCAL MAXIMUM RESIDUE LIMITS, BUT THE IMPORT TOLERANCES OF OTHER COUNTRIES MIGHT POSSIBLY BE EXCEEDED. IF THE CROP TO BE TREATED IS INTENDED FOR EXPORT, CONSULT THE RELEVANT EXPORTING ORGANISATION REGARDING WITHHOLDING PERIODS.

- Handle with care.
- Poisonous when absorbed through the skin, swallowed or inhaled.
- Toxic to animals and birds.
- Store in a cool, dry, well-ventilated place in the original container, tightly closed.
- Store away from food, feeds, seed, fertilizers and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Keep livestock and domestic animals out of treated areas for at least 24 hours.
- Re-entry: Do not enter treated field within 1 day after application unless wearing protective clothing.
- **In case of poisoning call a physician immediately and make this label available to him/her.**

Aerial application:

Notify all inhabitants in the immediate vicinity of the area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate water or adjacent areas.

Do not use aerial application in apples or pears.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest to the remedy concerned, as well as by the method, time and accuracy of the application. The registration holder furthermore, does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for the lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of uncertainty.

PRECAUTIONS

- Avoid inhalation of the spray mist or fumes.
- Avoid eye and skin contact.
- Wear a face mask and rubber gloves when handling, preparing and applying the spray mixture.

- Wash with soap and water after use or in the event of accidental skin contact.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke while mixing and applying the product, before washing hands and face and change of clothing.
- Prevent drift onto other crops, grazing, rivers, dams or any other area not under treatment.
- Clean the applicator thoroughly after use and dispose of the wash water where it will not contaminate food, grazing, rivers or dams.
- Containers and packages must be completely emptied before being disposed of. Bury in an approved, designated landfill.
- Destroy the container/bag by perforation and dispose of it in a safe way.
- **Never** re-use the empty container/bag for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

SYMPTOMS OF HUMAN POISONING

Pirimicarb is a carbamate compound, which inhibits cholinesterase. Systemic effects may begin within a few minutes or several hours of exposure. Initial symptoms can be nasal hyperaemia, watery discharge, chest discomfort and wheezing. Other symptoms may include nausea, vomiting, diarrhoea, abdominal cramps, headache, vertigo, tightness of the chest, anxiety, ocular pain, ciliary muscle spasm, blurred vision, small pupils, or mydriasis, lacrimation, salivation, sweating, and confusion. In severe cases, there may be involuntary defaecation, bradycardia, hypotension, pulmonary oedema, convulsions, coma and death from respiratory failure or cardiac arrest.

FIRST AID TREATMENT

- **Get medical attention in all cases.**
- **Skin:** Remove contaminated clothing and rinse the contaminated body area thoroughly with non-abrasive or mild detergent and large amounts of water, until no evidence of chemical remains. Do not rub the skin. Get medical attention if irritation occurs and persists.
- **Eyes:** Flush eyes with clean water for at least 20 minutes, occasionally lifting upper and lower lids. If irritation occurs and persists, obtain medical attention if necessary.
- **Inhalation:** Remove patient from the source of contamination and keep him/her calm and at rest. Keep air passage open. Get medical attention immediately.
- **Ingestion:** Get medical attention immediately. Do not give anything by mouth. If the person is alert, rinse mouth thoroughly with water.

NOTE TO PHYSICIAN

If consciousness level declines or vomiting has not occurred in 15 minutes empty stomach by gastric lavage. Avoid aminoglycosides and succinylcholine. Phenothiazines, reserpine, theophylline, morphine, Pralidoxime (2-PAM, Protopam) and other oximes are contra-indicated.

Antidote: Administer atropine sulphate intravenously or intramuscularly. In moderately severe poisoning administer atropine sulphate, 0.4 to 2.0 mg repeated every 15 minutes, until atropinization is achieved (dry, flushed skin, dry mouth and tachycardia). Maintain atropinization by repeated doses for 2 to 12 hours, or longer, depending on the severity of poisoning.

RESISTANCE WARNING

UNIPEX 500 WDG is a group code 1A insecticide. Any insect population may contain individuals naturally resistant to **UNIPEX 500 WDG** and other group code 1A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **UNIPEX 500 WDG** or any other group code 1A insecticide.

To delay insecticide resistance:

- avoid exclusive repeated use of insecticides from the same insecticide group code. Alternate or tank-mix with products from different insecticide group codes,
- integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

The full range of compatibility of **UNIPEX 500 WDG** with other remedies is not known; therefore, do not use it in tank mixtures with products other than those mentioned on this label.

Add a suitable adjuvant at the recommended label rates if necessary.

Mixing instructions:

- Fill the spray tank with clean water to just below the total volume required.
- Make a slurry of the **UNIPEX 500 WDG** to be added to the tank in a suitable container, ensuring that all the chemical has dispersed.
- Add the mixture to the water while stirring.
- Fill the spray tank with water to the required level while maintaining agitation, to ensure thorough mixing.
- Prepared spray mixtures must not be left in the spray tank for any length of time, e.g. overnight.

NOTE

UNIPEX 500 WDG does not affect most of the natural enemies which attack aphids on crops.

Where the crop foliage is dense, ensure that the spray also penetrates to the lower parts of the plant.

Aerial application:

Aerial application of **UNIPEX 500 WDG** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- Volume: A spray mixture volume of 30 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aurally at a lower volume rate than recommended above.
- Droplet coverage: 25 to 35 droplets per cm² must be recovered at the target area.
- Droplet size: A droplet spectrum with a VMD of 280 to 300 micron is recommended. Limit the production of fine droplets less than 150 micron (high drift and evaporation potential) to a minimum.
- Flying height: Maintain the height of the spray boom at 3 to 4 metre above the target. Do not spray when the aircraft dives, is in a climb or when banking.
- Use suitable atomising equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8 °C.
- Stop spraying if the wind speed exceeds 15 km per hour.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),
 - b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

APPLICATION RATES

Crop / Pest	Dosage Rate	Remarks
Apples Aphids	50 g / 100 ℓ water (5 g / 10 ℓ water)	Spray as soon as an infestation is noticed. Repeat when necessary.
Citrus Aphids	40 g / 100 ℓ water (4 g / 10 ℓ water)	Spray as and when necessary.
Cotton Aphids (except resistant strains of <i>Aphis gossypii</i>)	150 g / ha (7.5 g / 10 ℓ water)	Ground application: Apply in 200 litres water.
	175 g / ha	Aerial application: Apply in 30 litres water.
		Dryland cotton Spray when Aphids first appear. Irrigated cotton Apply when honeydew on plants is noticed. Ensure thorough coverage and repeat when necessary.
Cruciferae Cabbage aphid (<i>Brevicoryne brassicae</i>)	250 g (5 g / 10 ℓ water)	Apply in 500 litres water. Spray when Aphids first appear and repeat every 2 weeks. Direct sprays into the centre of the plants. Sprays will not control Aphids deep within the plants.
Groundnuts Aphids	250 g / ha (6 g / 10 ℓ water)	Apply in 400 litres water. Ensure thorough coverage.
Ornamentals Aphids	50 g / 100 ℓ water (5 g / 10 ℓ water)	Ensure thorough coverage of the plants to the point of run-off.
Peaches Peach aphid	50 g / 100 ℓ water (5 g / 10 ℓ water)	Ensure thorough coverage. Repeat application when re-infestation occurs.
Pecan nuts Aphids	100 g / 100 ℓ water (10 g / 10 ℓ water)	If Aphids are present, spray once in the middle of January.
Potatoes Aphids	500 g / ha (10 g / 10 ℓ water)	Apply in 500 litres water. Ensure thorough coverage.
Sorghum Honeydew aphid (<i>Melanaphis sacchari</i>) Funnel aphid (<i>Rhopalosiphum maydis</i>)	250 g / ha (17 g / 10 ℓ water)	Ground application: Apply in 150 litres water when the pest is increasing.
	250 g / ha	Aerial application: Apply in 30 litres water when the pest is increasing.
Wheat Green aphid (<i>Acythosiphon hordium</i>)	250 g / ha (17 g / 10 ℓ water)	Ground application: Apply in 150 litres water.
	250 g / ha	Aerial application: Apply in 30 litres water.
		NOTE Apply ONLY to wheat, which is over 200 mm in height.
Wheat and Oats Brown aphid (<i>Macrosiphum avenae</i>)	250 g / ha (17 g / 10 ℓ water)	Ground application: Apply in 150 litres water. Spray when the infestation of Aphids reaches an average of 5 Aphids per head.
	250 g / ha	Aerial application: Apply in 30 litres water. Spray when the infestation of Aphids reaches an average of 5 Aphids per head.