

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Fungicide



SULFLO 700 SC

Reg. No. L 6433 Act/Wet No. 36 of/van 1947

1: 26/4/2012 - May 2017

A suspension concentrate fungicide for the control powdery mildew (Oidium) on wine grapes.

'n Suspensie konsentraat swamdoder vir die beheer van poeieragtige skimmel (Oidium) op wyndruwe.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

sulphur 700 g/l swael

FRAC FUNGICIDE GROUP CODE **M2** FRAC SWAMDODER GROEPKODE



villa

Registration holder / Registrasiehouer:
Universal Crop Protection (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07
PO Box / Posbus 801, Kempton Park, 1620
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: 2902

Willow Set & Print 011 394-4486



CAUTION
VERSIGTIG



GEBRUIKSAANWYSINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED
REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

SULFLO 700 SC

Reg. No. L 6433 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: M2

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

sulphur / swael.....700 g/ ℓ

Registration holder / Registrasiehouer:

VILLA CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1992/002474/07 Mpy. Reg. Nr.

PO Box / Posbus 10413

ASTON MANOR, 1630

Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS**

- Harmful if inhaled or swallowed.
- Irritant of eyes, nose, throat and skin.
- Do not apply after the berries have reached pea size (4 to 6 weeks before harvest), as this may adversely affect fermentation.
- Do not apply when the temperature is above 32 °C.
- Do not apply within 4 weeks after budding.
- Store away from food, feeds, seed, fertilizers and other agricultural chemicals.
- Store in a cool, dry, well-ventilated place in the original container, tightly closed and secured.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Avoid inhalation of the spray mist
- Avoid contact of product with skin and eyes.
- Wear a face shield and rubber gloves when handling the concentrate and during mixing.
- Wash with soap and water after use or accidental skin contact.
- In case of eye contact, immediately flush the eyes with clean, cold water for 15 minutes, holding the eyelid(s) open. If irritation persists, seek medical advice.
- Wash contaminated clothes daily after use.
- Do not eat, drink or smoke whilst applying or mixing, or before washing hands and face or change of clothing.
- Prevent spray drift onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean the spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening and dispose of it in a safe way.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

DIRECTIONS FOR USE: Use only as directed.**Compatibility:**

SULFLO 700 SC is compatible with **Copper oxychloride 850 WP**, **Magnesium sulphate**, **Zinc oxide** and most organic pesticides.

Mixing instructions:

- Shake contents well before use by rocking/shaking a few times.
- Replace container cap after pouring.
- Always use spray tank strainer when pouring **SULFLO 700 SC** into spraying tanks.
- After each day's spraying flush out spray tank thoroughly with clean water.
- Use an automatic agitator in the tank when spraying.

Application:

- All treatments must be applied with suitable spray equipment that is in good working order and correctly calibrated to give the desired droplet size and coverage of the target area.

APPLICATION RATES

<i>Crop / Disease</i>	<i>Dosage</i>	<i>Remarks</i>
Vines <u>Wine grapes only:</u> Powdery mildew	1 per 100 litres water	Apply 500 to 1000 litres spray mixture per hectare depending upon development stage of vines. Start application when shoots are 10 to 15 cm long and repeat at 2 week intervals.

The following product mentioned in this label may be replaced with an equivalent product:

- **COPPER OXYCHLORIDE 850 WP** (L 6231) = **COPPER OXYCHLORIDE 850 WP** (L 5809).

COPPER OXYCHLORIDE 850 WP is a registered product of / is 'n geregistreerde produk van
Universal Crop Protection (Pty) Ltd.

COPPER OXYCHLORIDE 850 WP is a registered product of / is 'n geregistreerde produk van
Villa Crop Protection (Pty) Ltd.