

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Insecticide



ADVANCE 150 SC

Reg. No. L 9147 Act/Wet No. 36 of/van 1947
W 130689

5-10-3-2021 - March 2021

A suspension concentrate stomach and contact insecticide for the control of various insect pests as listed.

'n Suspensie konsentraat maag en kontak insekdoder vir die beheer van verskeie plae soos gelys.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

indoxacarb (oxadiazine) 150 g/l indoksakarb (oksadiasien)

IRAC INSECTICIDE GROUP CODE **22A** IRAC INSEKDODER GROEPKODE



villa

Registration holder / Registrasiehouer:
Universal Crop Protection (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07
PO Box / Posbus 801, Kempton Park, 1620
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: 3082



HARMFUL
SKADELIK

Willow Set & Print 011 394-4486



GEBRUIKSAANWYINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

DIRECTIONS FOR USE ENCLOSED

Batch number:
Lotnommer:

ADVANCE 150 SC

Reg. No. L 9147 / W 130689 Act/Wet No. 36 of/van 1947

IRAC INSECTICIDE GROUP CODE / IRAC INSEKODER GROEPKODE: 22A

ACTIVE INGREDIENT/ AKTIEWE BESTANDDEEL:

indoxacarb (oxadiazine) / indoksakarb (oksadiasien)..... 150 g/l

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

PO Box / Posbus 801,

KEMPTON PARK, 1620, Tel. (011) 396 2233

**HARMFUL****SKADELIK****WARNINGS****Withholding periods:**

Minimum time between the last application and harvest:	
Apples	28 days
Blueberries and Raspberries	42 days
Cruciferae, Cucurbits and Green Beans	3 days
Dry Beans (grazing)	42 days
Hops & Oats	7 days
Lettuce	1 day
Peas	7 days
Stone Fruit (Peaches and Nectarines)	28 days
Soybeans	14 days
Soybeans (grazing)	42 days
Canola	56 days
Maize and Sweetcorn (grazing)	42 days
Maize and Sweetcorn	7 days
Maize and Sweetcorn (tank mixtures with Tempes 100 EC – including grazing)	56 days
Maize (tank mixtures with Prime 50 EC or Methomyl 90 SP) – no grazing	42 days
Grain sorghum	56 days
Tobacco	56days
Cotton	14 days
Tomatoes	1 day
Strawberries	1 day
Sugarcane	28 days
Table grapes	14 days
Wine grapes	28 days
Peppers	1 day
Wheat, Barley & Oats (grazing)	42 days

WITHHOLDING PERIODS ARE BASED ON GOOD AGRICULTURAL PRACTICE AND WHEN FOLLOWED, LOCAL (SA) MRL'S WILL BE MET. FOR PRODUCE DESTINED FOR EXPORT, CONSULT YOUR EXPORT AGENT OR SUPPLIER FOR THE LATEST WITHHOLDING PERIODS RELEVANT TO YOUR EXPORT MARKET.

- Do not graze treated cotton.
- Handle with care.
- Keep out of reach of children, animals and uninformed persons.
- May be harmful if swallowed or inhaled.
- May cause an allergic skin reaction.
- Causes damage to organs (blood, nervous system and heart) through prolonged or repeated exposure.
- Toxic to animals, bees and fish.
- Toxic to aquatic life with long lasting effects.
- Store in a cool, dry place away from food, feeds, fertilizers and other agricultural remedies.
- Re-entry: Do not enter treated area within 1 day after treatment unless wearing protective clothing.
- **In case of poisoning call a physician and make this label available to him/her.**

Aerial application:

Notify all inhabitants of the immediate area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate water of adjacent areas.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation and the environment, or harm to man or animal, or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Do not inhale the spray mist.
- Wear protective clothing – suitable overalls, rubber boots, gloves and a face mask when preparing the spray mixture and during application.
- Avoid contact with eyes and prolonged or repeated skin contact.
- Remove and wash contaminated clothing after use.
- Wash with soap and water after use or accidental skin contact.
- Do not eat, drink or smoke whilst mixing, applying the product or before washing hands and face or change of clothing.
- Use only outdoors (mixing of the product included) and in a well-ventilated area.
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment by using a suitable drift retardant such as **INTERLOCK**[®] (L 10254 / W 130875 / N-AR 1856) in row crops or **MASTERLOCK**[®] (L 10496) in orchard crops.
- Thoroughly clean spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams. Clean the applicator with a household ammonia solution (1 %) or **Protank**[®] liquid cleaner before using other pesticides. Let the solution stand for several hours, preferably over-night. Rinse at least twice. This applicator should not be used for applying chemicals other than insecticides
- **TRIPLE RINSE THE EMPTY CONTAINER AS FOLLOWS:** Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.
- Prevent contamination of food, feeds, drinking water and eating utensils.

RESISTANCE WARNING

ADVANCE 150 SC is a group code 22A insecticide. Any insect population may contain individuals naturally resistant to **ADVANCE 150 SC** and other group code 22A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **ADVANCE 150 SC** or any other group code 22A insecticide.

To delay insecticide resistance:

- avoid exclusive repeated use of insecticides from the same insecticide group code. Alternate with products from different insecticide group codes,
- integrate other control methods (chemical, cultural, biological) into insect control programmes.
- **Do not** exceed the maximum number of applications per season with **ADVANCE 150 SC** (refer to instructions under “**USE RESTRICTIONS**” on this label).
- Avoid exposure of multiple successive pest generations to the same mode of action by making use of a window approach or block applications, consisting of 2 or 3 consecutive applications of **ADVANCE 150 SC**. A treatment window should not exceed the generation time of the pest (usually 60 days).
- Following a window of use with **ADVANCE 150 SC** or other group 22 insecticide, rotate to a window using a registered insecticide with a different mode of action.
- The period between group 22 treatment windows should not be less than 60 days from the last application with a group 22 insecticide.
- Monitor insect populations and apply **ADVANCE 150 SC** as per label instructions once locally determined economic thresholds have been reached. Multiple applications may be required to obtain effective control of an infestation.

For specific information on resistance management, contact the registration holder of this product or consult the website of the Insecticide Resistance Action Committee (<https://irac-online.org/>). If resistance is suspected, please consult your local company representative or agricultural advisor.

USE RESTRICTIONS

- Do not exceed the maximum number of applications of **ADVANCE 150 SC** as specified in the “**APPLICATION RATES**” table below for each crop-pest scenario. If the maximum number of applications are not specified, do not exceed two (2) applications of **ADVANCE 150 SC** per season.

DIRECTIONS FOR USE: Use only as directed.

General information & selectivity:

- Follow the label recommendations precisely for application rates, spray intervals and the optimal timing to apply **ADVANCE 150 SC**.
- **ADVANCE 150 SC** has minimal effect on beneficial insects and mites, thus promoting natural pest control.
- **ADVANCE 150 SC** is mainly a Lepidopteran (caterpillar type larvae of moths & butterflies) specific insecticide.
- **ADVANCE 150 SC** only controls larval (developmental) stages and not adult moths or butterflies.
- **Bees:** **ADVANCE 150 SC** can be dangerous to bees. To protect bees and other pollinators, do not apply **ADVANCE 150 SC** when honeybees are actively foraging.
- **Climatic conditions:** **ADVANCE 150 SC** is effective under hot climatic conditions. Once the spray mixture has dried on the target area, **ADVANCE 150 SC** will not wash off through rainfall or irrigation and these conditions will therefore not influence the normal residual activity of the product.

Mode of action:

- **ADVANCE 150 SC** acts by inhibition of sodium ion entry into nerve cells. This results in paralysis and death of the pest within 1 to 2 days. However, inhibition of the pest’s feeding occurs within 2 to 8 hours.
- **ADVANCE 150 SC** acts as a larvicide through ingestion (stomach action), as well as cuticular absorption (contact action).

Compatibility:

- **ADVANCE 150 SC** is compatible with the insecticides - **Lambda 50EC, Tempes 100 EC, Lambda Secure 106 CS, Baritone 50 EC, Methomyl 90 SP, Villa 51** and **Aquabuff Plus**, organosilicon surfactants like **Charge** and **Break-Thru® S240** and the drift retardant adjuvants **Interlock®** and **Masterlock®**.
- **Masterlock®** is an adjuvant for use with post-emergence foliar pesticide applications on permanent crops, orchards, and vineyards to reduce spray drift, improve droplet deposition, as well as canopy penetration and droplet spreading and absorption. Apart from buffers, **Masterlock®** should not be used with other adjuvants. Groups of pesticides/chemicals will behave in a similar way when applied with **Masterlock®** and as such, the effect can be extrapolated within a group. Consequently **Masterlock®** was not tested in all crop situations listed on this label and thus should not be applied where the pre-harvest interval is seven (7) days or less, unless specified.
- **Interlock®** can be added to improve drift control, canopy penetration and coverage in row crops. Similar to **Masterlock®**, **Interlock®** was also not tested in all crop situations listed on this label and thus should not be applied where the pre-harvest interval is seven (7) days or less, unless specified.
- The compatibility of **ADVANCE 150 SC** with other products may be influenced by several factors. As certain factors may vary from time to time, a physical compatibility test must always be performed before such a tank mixture is sprayed. If the products are physically compatible, this does **NOT** imply that they will be biologically compatible (effective against the target pest), thus the registration holder will not take responsibility for ineffective control in these scenarios.
- When **ADVANCE 150 SC** is used in conjunction with any other agricultural remedy, adhere to all **WARNINGS, PRECAUTIONS** and **DIRECTIONS FOR USE** mentioned on the relevant labels.

Mixing instructions:

- Half fill the spray tank with clean water. Add the required amount of **ADVANCE 150 SC** while maintaining agitation. Complete the filling operation whilst maintaining agitation.
- When mixing **ADVANCE 150 SC** with other products (as registered on this label), use the following procedure:
 - If a buffer is required, it should be added to the spray mixture first.
 - Mix the required quantity of WDG or WP products in a small quantity of water to make a smooth cream and add this to the spray mixture. If dry formulations are contained in water-soluble bags, add this to the spray mixture.

- In the following order, add any additional products: SC or SE formulations, followed by EC, EW or ME formulations. Lastly add products of SL, SG or SP formulation types. Pre-mix products individually before adding them to the spray mixture.
- After all the products have been thoroughly incorporated, add the wetter / spreader adjuvants such as **Charge, Villa 51, Interlock® or Masterlock®**.
- Maintain continuous suspension in the spray tank during mixing and application by means of proper agitation.
- The prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

Application instructions:

Always use suitable, correctly calibrated equipment, in good working order, to ensure thorough wetting of all parts of the plant.

Ground Application:

ADVANCE 150 SC can be applied with most generally used, purposed designed spraying equipment used in agriculture that meets above-mentioned requirements.

ADVANCE 150 SC has not been tested at low volume rates (<100 litres per hectare), therefore the registration holder cannot guarantee efficacy of use under such conditions.

Aerial application:

Aerial application of **ADVANCE 150 SC** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- Volume: A spray mixture volume of 30 to 35 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aurally at a lower volume rate than recommended above.
- Droplet coverage: 35 to 45 droplets per cm² must be recovered at the target area.
- Droplet size: A droplet spectrum with a VMD of 250 to 280 micron is recommended. Limit the production of fine droplets less than 150 micron (high drift and evaporation potential) to a minimum.
- Flying height: Maintain the height of the spray boom at 3 to 4 metres above the target. Do not spray when aircraft dives, is in a climb or when banking.
- Use suitable atomizing equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomizers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8°C.
- Stop spraying if the wind speed exceeds 15 km per hour.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - a) Reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),
 - b) Damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

Application through Irrigation systems:

Ensure that the Pivot irrigation system complies with the following specifications, before application starts:

- The distribution coefficient of the pivot irrigation system must be >90 % and the Pivot should not apply more than 5 mm (50 000 litres) per hectare at maximum speed.
- Clean all the sieves of the pivot irrigation system before starting application. Calibrate the injection pump of the pivot irrigation system according to the time it takes for the pivot irrigation system to complete one full circle.

- Set the pivot irrigation system to maximum speed so that as little water as possible (not more than 5 mm) is applied per hectare.
- Mix **ADVANCE 150 SC** and a mineral oil adjuvant. Stir the spray mixture continuously before and during application.
- Inject the **ADVANCE 150 SC** spray mixture in the main application line of the pivot irrigation system as soon as the pivot irrigation system reached maximum speed. Mark the position on the land when the spray mixture reaches the furthest end of the pivot irrigation system.
- Complete one full circle from the aforementioned point.

APPLICATION RATES

Crop / Pest	Dosage rate	Remarks
<u>Apples</u> <u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)	50 ml / 100 ℓ water (5 ml / 10 ℓ water)	<u>Foliar Application:</u> Apply as a full cover spray in 500 to 2000 litres water per hectare. Apply preventively or when eggs or larvae are present, but before larvae enter the fruit. A follow-up application may be necessary 10 to 14 days later depending on re-infestation of the pest. Thorough coverage is essential. Do not exceed two (2) ADVANCE 150 SC applications per season.
<u>Blueberries and Raspberries</u> <u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)	40 ml / 100 ℓ water (4 ml / 10 ℓ water)	Apply preventively as a foliar spray as soon as bollworm eggs or small larvae are present as confirmed by regular scouting of the crop. Apply as a full cover spray in 500 to 1200 litres water per hectare depending on plant size. If required, apply a follow-up treatment 10 to 14 days later (e.g., under conditions of continuous re-infestation). Ensure thorough and even coverage of the foliage. Caterpillars that are obscured by dense foliage may not be controlled effectively. Refer to “IMPORTANT NOTES” below.
<u>IMPORTANT NOTES WITH REGARDS TO BLUEBERRIES AND RASPBERRIES</u>		
<ul style="list-style-type: none"> • Do not exceed two (2) ADVANCE 150 SC applications per season. Should further applications be necessary, use a product with a different mode of action. • Where the fruit of these crops is destined for the export market, it is recommended to apply ADVANCE 150 SC only before flowering or after harvesting of the fruit (i.e., no fruit present on the trees). Should control of the pest be required during the fruiting period, apply a different registered pesticide. • The addition of a non-ionic adjuvant (e.g., Villa 51) to the spray mixture is recommended to improve coverage. 		
<u>Strawberries</u> Larvae of African bollworm (<i>Helicoverpa armigera</i>)	30 ml / 100 ℓ water PLUS 20 ml / 100 ℓ water Charge	Apply as a corrective foliar application as soon as the first bollworm eggs or small larvae are observed during regular scouting. Apply as a full cover spray ensuring good penetration and coverage of all plant parts. A follow-up application can be made after 10 days. Do not exceed two (2) applications of ADVANCE 150 SC per growing season. If further control is needed, alternate with chemicals with a different mode of action.

Crop / Pest	Dosage rate	Remarks
Cotton	VERY IMPORTANT	Refer to “USE RESTRICTIONS”, “DIRECTIONS FOR USE” (including “Crop information”) and “APPLICATION INSTRUCTIONS”.
Larvae of: African bollworm (<i>Helicoverpa armigera</i>) Red bollworm (<i>Diparopsis castanea</i>) Spiny bollworms (<i>Earias biplaga</i> & <i>E. insulana</i>)	<u>Broadcast ground application:</u> 200 to 250 ml / ha (5 to 6.5 ml / 10 l water) ADVANCE 150 SC PLUS Aquabuff Plus	
Nymphs of: Cotton leafhoppers (<i>Jacobiella fascialis</i> & <i>J. libyca</i>)	<u>Aerial application:</u> 250 ml / ha ADVANCE 150 SC PLUS 1 l / 100 litres water (minimum 350 ml / ha) mineral oil adjuvant	Spray volumes are based on plants taller than 0.6 m. Apply in 200 to 400 litres spray mixture per hectare and ensure thorough foliage cover. Preventive treatment: Use the lower rate in a weekly preventive programme or when economic threshold values are reached. Refer “NOTES ON ECONOMIC THRESHOLD LEVELS” below. Corrective treatment: Use the higher application rate for corrective application when the economic threshold value is exceeded. Refer “NOTES ON ECONOMIC THRESHOLD LEVELS” below. Apply a minimum of 35 litres spray mixture per hectare. Ensure thorough coverage of the foliage.
	<ul style="list-style-type: none"> • ADVANCE 150 SC can be used in a 7 to 14-day interval. Use the shorter interval when plants are actively growing or when leafhoppers are present. • If infestation persists, use a registered product with a different mode of action to provide further control. Do not exceed two (2) applications of ADVANCE 150 SC per season. • ADVANCE 150 SC will control all larval instars of the African bollworm. • Most beneficial insects and mites are not affected by ADVANCE 150 SC. It is therefore recommended to use ADVANCE 150 SC early in the season of the spray programme. 	

NOTES ON ECONOMIC THRESHOLD LEVELS

- Bollworm species**
About 8 to 20 weeks after emergence, start scouting by inspecting 24 plants per 15 hectare and base spray decisions thereon.
- African bollworm**
Egg threshold: 12 per 24 plants
Larval threshold: 5 per 24 plants
- Red bollworm**
Egg threshold: 6 per 24 plants
Larval threshold: 2 per 24 plants
- Spiny bollworms**
Larval threshold: 2 per 24 plants
- Leafhoppers**
Start application when 2 Leafhopper adults/nymphs are present per leaf.

Crop / Pest	Dosage rate	Remarks
<p>Maize & Sweetcorn</p> <p>Cutworm (<i>Agrotis</i> spp.)</p>	<p>3 ml / 100 m row (10 ml / 10 l water)</p>	<p>Apply as a preventive (pre-emergence) application in at least 3 liters water per 100 m row length.</p> <p>Apply as a row treatment in a 30 cm wide band over the row just after planting. The soil to be treated must be well prepared – free of clods and excessive plant debris.</p> <p>Apply only if the top 3 cm of the soil is moist.</p>
<p>Larvae of: Maize stalk borer (<i>Busseola fusca</i>)</p> <p>Sorghum stem borer (<i>Chilo partellus</i>)</p> <p>African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>VERY IMPORTANT Refer to “USE RESTRICTIONS”, “DIRECTIONS FOR USE” (including “Crop information”) and “APPLICATION INSTRUCTIONS”.</p> <p>300 ml / ha (10 ml / 10 l water)</p> <p>ADVANCE 150 SC</p> <p>PLUS</p> <p>120 ml / ha (4 ml / 10 l water)</p> <p>Lambda 50EC</p> <p>OR</p> <p>60 ml / ha (2 ml / 10 l water)</p> <p>Lambda Secure 106 CS</p> <p>PLUS</p> <p>Aquabuff Plus</p> <hr/> <p>3 ml / 100 m plant row ADVANCE 150 SC</p> <p>PLUS</p> <p>1.2 ml / 100 m plant row Lambda 50EC</p> <p>OR</p> <p>0.6 ml / 100 m plant row Lambda Secure 106 CS</p> <p>PLUS</p> <p>Aquabuff Plus</p>	<p>Broadcast ground application: Apply as a preventive or early corrective treatment. Apply 300 to 450 litres of spray mixture per hectare. Apply over the funnel of the plants and ensure thorough coverage of the foliage. To ensure optimum results, application must take place early in the morning when dew is present. The absence of rain within 3 days of application or irrigation after application can lead to a decrease in control with ADVANCE 150 SC plus Lambda 50EC OR Lambda Secure 106 CS spray mixtures. Water after application is essential to wash the spray mixture in the funnel.</p> <p>Under conditions of repeated infestation, apply ADVANCE 150 SC plus Lambda 50EC OR Lambda Secure 106 CS in a 10 to 14-day spray programme. Use the shorter spray interval when the pest infestation pressure is high or when maize is growing fast.</p> <hr/> <p>Aerial application: As above in at least 30 litres water per hectare.</p> <hr/> <p>Ground directed row application: Apply 3 litres spray mixture per 100 m plant row over the funnel of the plants and ensure thorough coverage of the foliage. To ensure optimum results, application must take place early in the morning when dew is present.</p> <p>The absence of rain within 3-days of application or irrigation after application can lead to a decrease in control with ADVANCE 150 SC plus Lambda 50EC OR Lambda Secure 106 CS spray mixtures. Water after application is essential to wash the spray mixture into the funnel.</p> <p>Under conditions of repeated infestation, apply ADVANCE 150 SC plus Lambda 50EC OR Lambda Secure 106 CS in a 10 to 14-day spray programme. Use the shorter spray interval when the pest infestation pressure is high or when maize is growing fast.</p>

Crop / Pest	Dosage rate	Remarks
<p><u>Maize & Sweetcorn (continued)</u></p> <p><u>Larvae of:</u> Maize stalk borer (<i>Busseola fusca</i>)</p> <p>Sorghum stem borer (<i>Chilo partellus</i>)</p> <p>African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Application through pivot irrigation system:</u></p> <p>300 ml / ha ADVANCE 150 SC</p> <p>PLUS</p> <p>120 ml / ha Lambda 50EC</p> <p>OR</p> <p>60 ml / ha Lambda Secure 106 CS</p> <p>PLUS</p> <p>5 l / ha mineral oil adjuvant</p>	<p>IMPORTANT Refer to “Application through irrigation systems” above. Apply as a preventative or early corrective treatment. Under conditions of repeated infestation, apply ADVANCE 150 SC plus Lambda 50EC OR Lambda Secure 106 CS in a 10 to 14-day spray programme. Use the shorter spray interval when the pest infestation pressure is high or when maize is growing fast.</p>
<p><u>Larvae of:</u> Maize stalk borer (<i>Busseola fusca</i>)</p> <p>Sorghum stem borer (<i>Chilo partellus</i>)</p>	<p><u>Broadcast ground application:</u></p> <p>300 ml / ha (10 ml / 10 l water) ADVANCE 150 SC</p> <p>PLUS</p> <p>300 ml / ha (10 ml / 10 l water) Tempes 100 EC</p> <p>PLUS</p> <p>25 ml / 100 l water (2.5 ml / 10 l water)</p> <p>Charge</p>	<p>Apply as a preventive or early corrective treatment. Apply 300 to 450 litres of spray mixture per hectare. Apply over the funnel of the plants and ensure thorough coverage of the foliage. To ensure optimum results, application must take place early in the morning when dew is present. The absence of rain within three (3) days following application or irrigation after application can lead to a decrease in control with ADVANCE 150 SC plus Tempes 100 EC spray mixtures. Water following application is essential to wash the spray mixture into the funnel.</p> <p>Under conditions of repeated infestation, apply a second ADVANCE 150 SC plus Tempes 100 EC treatment 7 to 10-days later. Use the shorter spray interval when the pest infestation pressure is high or when maize is growing fast.</p>

Crop / Pest	Dosage rate	Remarks
Maize Fall armyworm (<i>Spodoptera frugiperda</i>)	3 ml / 100 m row PLUS 7 ml / 100 m row Baritone 50 EC OR 280 to 560 g / ha Methomyl 90 SP PLUS 25 ml / 100 l water Charge	Apply in 450 litres of water, at the beginning of infestation, and at the first signs of damage. For optimal results, applications must be made early in the morning when dew is present. Ensure thorough coverage of all plant parts. Application is not recommended for the control of caterpillars larger than 1 cm. Apply in a block application with two (2) consecutive applications at a 7-day interval. Do not exceed two applications of ADVANCE 150 SC per season. If further control is needed, alternate with products with a different mode of action.
<u>IMPORTANT NOTES ON STALK BORER, STEM BORER, AFRICAN BOLLWORM AND FALL ARMYWORM CONTROL ON MAIZE</u>		
<ul style="list-style-type: none"> • Do not apply ADVANCE 150 SC on maize that is under drought stress. • Larvae that have already tunneled into the funnels will not be controlled with ADVANCE 150 SC spray mixtures. • Do not exceed two (2) applications of ADVANCE 150 SC per season. Two (2) consecutive applications can be made whereafter a registered insecticide with a different mode of action should be used. • Maize stalk borer, <i>Busseola fusca</i>, larvae control: <ul style="list-style-type: none"> - Commercial maize: Apply preventively or when eggs are found on 5 % of the plants, or when 10 % of the plants are showing shot hole damage symptoms on the maize funnels, which are caused by small feeding larvae. - Sweetcorn: Apply preventively, or when egg laying starts, or with the first signs of shot hole damage symptoms on the maize funnels. - Applications must be made before larvae migrate to the stem of the maize of sweetcorn plant and when larvae are smaller than the second instar. Do not apply ADVANCE 150 SC against maize stalk borer while the tassels are encircled by the flag leaf. • Sorghum stem borer, <i>Chilo partellus</i>, larvae control: <ul style="list-style-type: none"> - Commercial maize: Apply preventively, or when eggs are found on 2.5 % of the plants, or when 5 % of the plants are showing shot hole damage symptoms on the maize funnels, which are caused by small feeding larvae. - Sweetcorn: Apply preventively, or when egg laying starts or with the first signs of shot hole damage symptoms on the sweetcorn funnels. - Applications must be made before larvae migrate to the stem of the maize of sweetcorn plant and when larvae are smaller than the second instar. Do not apply ADVANCE 150 SC against sorghum stem borer while the tassels are encircled by the flag leaf. • African bollworm, <i>Helicoverpa armigera</i>, larvae control: <ul style="list-style-type: none"> - Funnel infestation Apply as for maize stalk borer. - Cob infestation Apply when the first larvae are noticed on the beard during cob formation. Larvae that are already inside the beard or have migrated into the cobs will not be controlled. • Fall Armyworm, <i>Spodoptera frugiperda</i>, larval control: <ul style="list-style-type: none"> - Commence application at the first signs of infestation when no more than 5 % of the plants show damage. - Applications are NOT recommended for control of larvae larger than 1 cm. - Applications must be done before the larvae migrate into the cobs of the maize plants. Larvae that are already deep within the beard or have migrated into the cobs will not be controlled. 		

Crop / Pest	Dosage rate	Remarks
<p><u>Sugarcane</u></p> <p>Larvae of: Sugarcane borer (<i>Eldana saccharina</i>)</p>	<p>Ground application: 300 ml / ha</p> <p>Aerial application: 300 ml / ha</p>	<p>VERY IMPORTANT Refer to “USE RESTRICTIONS”, “DIRECTIONS FOR USE” (including “Crop information”) and “APPLICATION INSTRUCTIONS”.</p> <p>Early Corrective Application: <u>Ground application:</u> Apply in 350 to 500 litres water per hectare.</p> <p><u>Aerial application:</u> Apply in at least 30 litres water per hectare.</p> <p>Scouting of fields are important to determine Sugarcane borer control strategies. The Sugarcane borer larvae need to feed on the treated foliage in order to be controlled. Thus, optimal control with ADVANCE 150 SC depends on a full cover application to the foliage of the crop. Larvae feeding on unsprayed foliage will not be controlled. Although ADVANCE 150 SC will control larvae of all stages of development, including the large 5th instar larvae, larvae that have penetrated the stalks or are obscured by dense foliage during application may not be adequately controlled. Therefore, it is essential to ensure thorough coverage of the foliage.</p> <p>Do not exceed four (4) applications of ADVANCE 150 SC per season. Applications can be made at a one-month interval. If more treatments are required, use registered insecticides from another IRAC mode of action group. Most beneficial insects and predatory mites are unaffected by applications of ADVANCE 150 SC.</p>
<p><u>NOTES</u></p> <ul style="list-style-type: none"> • The use of ADVANCE 150 SC is only recommended when plants are growing actively. • Do not apply ADVANCE 150 SC on sugarcane that is under drought stress. Larvae that have already tunnelled into the stalks will not be controlled by ADVANCE 150 SC spray mixtures. • Do not irrigate within six (6) hours after an ADVANCE 150 SC application or apply the product when rain is expected within six (6) hours after application, to allow spray mixture to dry. Once the spray mixture has dried it will not easily be washed off by rain or irrigation. <p>Using “Cultural Control Practices” to assist with control of Sugarcane borer (<i>Eldana saccharina</i>) larvae in sugarcane.</p> <ul style="list-style-type: none"> • Plant clean, non-infested seed cane in order to improve crop stands. • Early harvesting helps to reduce Sugarcane borer numbers, as Sugarcane borer numbers and damage accumulate as the crop ages, especially after approximately 12 months. • Crops growing under stress conditions are more liable to attack. Under extreme conditions (e.g. drought) damage may increase significantly. • Burning heavily infested crops followed by good field hygiene may reduce the initial infestation. • Ensure that the stalks are cut as low as possible to ground level when harvesting. Above-ground stalk material may harbour larvae. • The “push-pull” sugarcane habitat management strategy is an option for biological control. • Choose cultivars according to the sugarcane borer risk in your area as differences exist between cultivar resistance to sugarcane borer. 		

Crop / Pest	Dosage rate	Remarks
<p><u>Canola</u></p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>250 to 300 ml / ha</p> <p>PLUS</p> <p>an organosilicon surfactant (such as Charge)</p>	<p>Apply preventatively/correctively as required in 300 to 600 litres water per hectare and ensure thorough coverage. The use of hollow or full cone nozzles is recommended. Use the higher dosage rate when applied correctively. Use the lower rate for subsequent applications when applied in a regular programme.</p> <p>Apply a block application of 2 to 3 consecutive ADVANCE 150 SC applications, and then switch to products with other modes of action.</p> <p>The addition of an organosilicon surfactant (such as Charge) is recommended to improve coverage.</p> <p>ADVANCE 150 SC controls all larval instars on contact.</p> <p><u>Aerial application:</u> Apply in at least 30 litres water per hectare.</p>
<p><u>Larvae of:</u> Diamond back moth (<i>Plutella xylostella</i>)</p>	<p>250 ml / ha</p> <p>PLUS</p> <p>an organosilicon surfactant (such as Charge)</p>	<p>Start application as soon as an infestation is noticed on leaves and flowers.</p> <p><u>Ground application:</u> Apply in 250 liters water per hectare. Ensure good coverage. Single application.</p> <p><u>Aerial application:</u> Apply in at least 30 litres water per hectare.</p>
<p><u>Cruciferae</u> (<u>Cabbage, Broccoli, Cauliflower and Brussel Sprouts</u>)</p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p> <p>Diamondback moth (<i>Plutella xylostella</i>)</p> <p>Cabbage webworm (<i>Hellula undalis</i>)</p> <p>Cabbage white butterfly (<i>Pieris brassicae</i>)</p>	<p><u>Ground Application:</u> 250 to 300 ml /ha (5 to 10 ml / 10 l water) (maximum 300 ml / ha)</p>	<p><u>Early Corrective Application:</u> Apply in 300 to 600 litres of water per hectare and ensure thorough coverage of the head, where the larvae feed. Make use of hollow or full cone nozzles to improve coverage within the structure of the crop. The higher application rate should be used when making a corrective application. The lower rate can be used for subsequent applications when applied as part of a regular spray programme.</p> <p><u>Diamondback moth and cabbage webworm</u> Application should commence shortly after transplant, before or as soon as the first eggs or larvae appear. Damage is worst at the early crop stage. Under conditions of continuous reinfestation, apply in a spray programme at 7 to 10-day intervals, using the shorter spray interval early in the season when crops are growing actively.</p> <p>Do not exceed three (3) applications of ADVANCE 150 SC per season. If further control is needed, alternate with chemicals with a different mode of action.</p> <p><u>Bollworm and Cabbage white butterfly</u> Apply correctively as needed.</p> <p>The alternation of ADVANCE 150 SC with products having a different mode of action is recommended. Apply a block application of 2 to 3 consecutive ADVANCE 150 SC applications, and then switch to products with different modes of action.</p> <p>Do not apply more than three (3) sprays of ADVANCE 150 SC per season, including when applied against different pest species.</p> <p>Most beneficial insects and predatory mites are unaffected by applications of ADVANCE 150 SC.</p> <p>The addition of a suitable registered adjuvant (e.g., Villa 51 or Charge) at the registered rate, is recommended to improve coverage.</p>

Crop / Pest	Dosage rate	Remarks
<p>Cucurbits (Pumpkins, Squash, Watermelons, Muskmelons, Baby Marrows, Patty Pans and Cucumbers)</p> <p>Larvae of: African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Ground Application:</u> 250 ml /ha (3.5 to 10 ml / 10 l water) (maximum 250 ml / ha)</p>	<p><u>Early Corrective Application:</u> Apply in 250 to 750 litres water per hectare depending on the specific cucurbit species and crop stage. Ensure thorough even coverage of the foliage. Apply preventively at flowering or when the first bollworm eggs or small larvae (less than 10 mm in length) are observed following regular scouting during flowering / fruit-set stages. It is important to take into consideration that even slight bollworm feeding damage on the flowers of cucurbit crops may result in significant yield losses. A follow-up application 7 to 10 days later will normally be necessary if the first application is made at the early flowering stage or under conditions of continuous re-infestation.</p> <p>Larvae that have penetrated the fruits, or are obscured by dense foliage, during application may not be adequately controlled. It is therefore essential to ensure thorough coverage of the foliage.</p> <p>Do not exceed two (2) applications of ADVANCE 150 SC per season. Should a third application be needed, use a product with a different mode of action. The addition of a suitable registered adjuvant (e.g. Villa 51 or Charge) at the registered rate, is recommended to improve coverage.</p>
<p>Drybeans</p> <p>Larvae of: African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>250 ml /ha (8.5 ml / 10 l water) (maximum 250 ml / ha)</p> <p>PLUS</p> <p>20 ml / 100 l water (2 ml / 10 l water)</p> <p>Charge</p>	<p><u>Corrective Ground Application:</u> Apply in 330 to 500 litres water per hectare and ensure thorough, even coverage of the foliage. <u>Aerial application:</u> Apply in at least 30 litres water per hectare. Ensure thorough, even coverage of the foliage.</p> <p>Apply at flowering or when the first bollworm eggs or small larvae (less than 10 mm in length) are observed following regular scouting during flowering and pod-set stages. A second application with an insecticide with a different mode of action could be necessary if the ADVANCE 150 SC application is made at the early flowering stage or under conditions of continuous re-infestation.</p> <p>Larvae that have penetrated the pods or are obscured by dense foliage during application may not be adequately controlled. It is therefore essential to ensure thorough coverage of the foliage.</p> <p>Do not exceed one (1) application of ADVANCE 150 SC per season in drybeans.</p>

Crop / Pest	Dosage rate	Remarks
<p><u>Green Beans</u></p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>250 ml / ha (7 to 10 ml / 10 l water) (maximum 250 ml / ha)</p>	<p><u>Ground Foliar Application:</u> Apply in 250 to 350 litres water per hectare and ensure thorough even coverage of the foliage. Apply preventively at flowering or when the first bollworm eggs or small larvae (less than 10 mm in length) are observed following regular scouting during flowering / fruit-set stages. A follow-up application 7 to 10 days later will normally be necessary if the first application is made at the early flowering stage or under conditions of continuous re-infestation.</p> <p>Larvae that have penetrated the pods or are obscured by dense foliage during application may not be adequately controlled. It is therefore essential to ensure thorough coverage of the foliage. The addition of a non-ionic adjuvant (e.g., Villa 51) to the spray mixture is recommended to improve coverage. Do not exceed two (2) applications of ADVANCE 150 SC per season. Should a third application be needed, use a product from a different IRAC mode of action.</p>
<p><u>Hops</u></p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Ground Application:</u> 30 ml / 100 l water</p>	<p>Only for use by authorized members of the SA Hop Growers Association and for production of hops for use in South Africa only.</p> <p><u>Early Corrective Foliar Application:</u> Apply in 500 to 1000 litres of water per hectare depending on plant height and density. Good coverage of all foliage is essential. Apply when the first larvae are observed usually early in the season on young shoots, during flowering and after flowering when cones are formed. Regular scouting of hop fields is essential to determine the timing of the first and subsequent applications if necessary. Apply ADVANCE 150 SC at a spray interval of 10 to 14 days. Use the shorter interval early in the growth season when plants are growing actively or under high pest pressure. Do not exceed two (2) applications per season with ADVANCE 150 SC. The alternation of ADVANCE 150 SC with insecticides that have a different mode of action is recommended. Apply a block application of two (2) consecutive ADVANCE 150 SC applications, and then switch to products with different modes of action. The addition of a non-ionic adjuvant (e.g., Villa 51) to the spray mixture is recommended to improve coverage.</p>
<p><u>Lettuce</u></p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>250 ml / ha (4 to 5 ml / 10 l water) (maximum 250 ml / ha)</p>	<p><u>Ground Application:</u> Commence with treatment in the initial phase of the crop (before the formation of the head), when the first larvae smaller than 10 mm are found in the crop. Apply in 500 to 600 litres of water per hectare depending on plant size. Do not exceed two (2) applications per season 7-days apart. The alternation of ADVANCE 150 SC with insecticides with a different mode of action is recommended. The addition of a suitable registered adjuvant (e.g., Villa 51 or Charge) at the registered rate, is recommended to improve coverage.</p>

Crop / Pest	Dosage rate	Remarks
<p>Peas (Green peas, Including “Mange Tout”)</p> <p>Larvae of: African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Ground Application:</u> 250 ml / ha (7 to 10 ml / 10 l water) (maximum 250 ml / ha)</p> <p><u>Aerial Application:</u> 300 ml / ha</p>	<p><u>Ground application:</u> Apply 250 to 350 litres spray mixture per hectare and ensure thorough even coverage of the foliage.</p> <p><u>Aerial application:</u> Apply in at least 30 litres water per hectare. Ensure thorough, even coverage of the foliage.</p> <p>Apply at flowering or when the first bollworm eggs or small larvae (less than 10 mm in length) are observed following regular scouting during flowering / fruit set stages. A follow-up application 7 to 10 days later will normally be necessary if the first application is made at the early flowering stage or under conditions of continuous re-infestation.</p> <p>Larvae that have penetrated the pods or are obscured by dense foliage during application may not be adequately controlled. It is therefore essential to ensure thorough coverage of the foliage.</p> <p>Do not exceed two (2) applications of ADVANCE 150 SC per season. Should a third application be needed, switch to a product from a different IRAC mode of action group.</p> <p>The addition of a non-ionic adjuvant (e.g., Villa 51) to the spray mixture is recommended to improve coverage.</p>
<p>Peppers</p> <p>Larvae of: African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>30 ml / 100 l water (3 ml / 10 l water)</p> <p>PLUS</p> <p>20 ml / 100 l water (2 ml / 10 l water)</p> <p>Charge</p>	<p>Regular scouting of pepper fields is essential to determine the timing of the first and subsequent application if necessary.</p> <p>Apply in 500 to 1500 litres spray mixture per hectare and ensure good coverage of all foliage. In the case of trellised peppers, both sides of the pepper row must be sprayed.</p> <p>Apply a second treatment 7 days later if necessary.</p> <p>Do not exceed two (2) applications of ADVANCE 150 SC per season.</p> <p>The addition of an organosilicon surfactant (such as Charge) is recommended to improve coverage.</p>
<p>Tomato leafminer (<i>Tuta absoluta</i>)</p>	<p><u>Ground application:</u> 300 ml / ha (when applying > 1000 l spray mixture per ha, use 30 ml / 100 l water)</p>	<p><u>Early corrective foliar application:</u> Monitor adult moth flights and commence application at egg-hatching or when the very first signs of the pest are noticed. Apply in 500 to 1500 litre spray mixture per hectare, ensuring thorough coverage of all plant parts.</p> <p>The addition of a suitable registered adjuvant is recommended to improve coverage e.g. Villa 51 at 50 ml per 100 litres or Charge at 20 ml per 100 litres of spray mixture.</p>
<p>NOTES ON PEPPERS:</p> <ul style="list-style-type: none"> Regular scouting is essential to determine optimal timing of the first and subsequent applications. In the case of trellised peppers, both sides of the row must be sprayed. Do not exceed four (4) applications of ADVANCE 150 SC per season. Apply ADVANCE 150 SC in block applications consisting of two (2) consecutive applications of ADVANCE 150 SC before switching to insecticides with a different mode of action. Applications should be made at 7-day intervals. Most beneficial insects are unaffected by applications of ADVANCE 150 SC. 		

Crop / Pest	Dosage rate	Remarks
<p><u>Soybeans</u></p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p> <p>Cabbage Semi-Looper (<i>Thysanoplusia orichalcea</i>)</p>	<p><u>Ground Application:</u> 250 ml / ha</p> <p>PLUS</p> <p>20 ml / 100 l water Charge</p> <p><u>Aerial Application:</u> 300 ml / ha</p>	<p><u>Ground application:</u> Apply in 300 to 700 litres water per hectare in soybeans. Ensure thorough, even coverage of the foliage.</p> <p><u>Aerial application:</u> Apply in at least 30 litres water per hectare. Ensure thorough, even coverage of the foliage.</p> <p>Apply as a corrective foliar application at flowering or when the first bollworm eggs or small larvae (less than 10 mm in length) are observed following regular scouting during flowering and pod-set stages. A follow-up application 7 to 10 days later will normally be necessary if the first application is made at the early flowering stage or under conditions of continuous re-infestation.</p> <p>Larvae that have penetrated the pods or are obscured by dense foliage during application may not be adequately controlled. It is therefore essential to ensure thorough coverage of the foliage.</p> <p>Do not exceed two (2) applications of ADVANCE 150 SC per season. Should a third application be needed, use an insecticide with a different mode of action.</p> <p>Most beneficial insects and predatory mites are unaffected by applications of ADVANCE 150 SC.</p> <p>The addition of a suitable registered adjuvant, at the registered rate, is recommended to enhance efficacy.</p> <p>Apply as a full cover spray when the pest is first noticed. The spray volume depends on the plant size (200 to 300 litres water per hectare).</p> <p>Apply a second treatment 7 days later if necessary.</p>
<p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Application through pivot irrigation system:</u></p> <p>250 ml / ha</p> <p>PLUS</p> <p>5 l / ha mineral oil adjuvant</p>	<p><u>IMPORTANT</u> Refer to “Application through irrigation systems” notes above.</p> <p>Apply as a preventative or early corrective treatment. Apply a second ADVANCE 150 SC treatment 7 days later.</p>
<p><u>Stone Fruit</u> (Peaches and Nectarines)</p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>30 ml / 100 l water (3 ml / 10 l water)</p>	<p><u>Foliar Application:</u> Apply as a full cover spray 500 to 2000 litres water per hectare and ensure thorough coverage. Apply preventively or when eggs or larvae are present, but before larvae enter the fruit.</p> <p>If required, apply a follow-up treatment 10 to 14 days later depending on infestation pressure of the pest.</p> <p>Do not exceed two (2) applications in total per season.</p>

Crop / Pest	Dosage rate	Remarks
<p>Table and Wine Grapes</p> <p>Larvae of: African bollworm (<i>Helicoverpa armigera</i>)</p>	<p>30 ml / 100 l water (3 ml / 10 l water)</p>	<p>Apply preventively, or as soon as eggs or larvae are present, but before larvae enter the fruit. If required, apply a follow-up treatment 10 to 14 days later depending on infestation pressure of the pest. Refer to “IMPORTANT NOTES” below.</p>
<p>Larvae of: False codling moth <i>Thaumatotibia leucotreta</i></p>	<p>40 ml / 100 l water (4 ml / 10 l water)</p>	<p>Always apply ADVANCE 150 SC as part of an Integrated Pest Management program (IPM). Commence with ADVANCE 150 SC applications when moth catches in pheromone traps confirms an infestation of False codling moth. Apply follow up treatments with 10-day intervals if required. For best control results, combine ADVANCE 150 SC applications with, or precede with other methods of False codling moth control such as mating disruption, biological control programs, good sanitation programs and alternative registered insecticides. Refer to “IMPORTANT NOTES” below.</p>
<p>IMPORTANT NOTES WITH REGARDS TO TABLE AND WINE GRAPES</p> <ul style="list-style-type: none"> • Apply as a full cover spray in 1000 to 1500 litres water per hectare depending on vine size and ensure thorough, even coverage of the plant structure. • Do not exceed two (2) ADVANCE 150 SC applications in total per season on crop. Should any further control of the pest be required use a product from a different IRAC mode of action group. • BEE SAFETY: According to the standards of Good Agricultural Practices, ADVANCE 150 SC should not be applied if honeybees are active. 		
<p>Tomatoes</p> <p>Larvae of: African bollworm (<i>Helicoverpa armigera</i>)</p> <p>Potato tuber moth (<i>Phthorimaea operculella</i>)</p> <p>Tomato semi-looper (<i>Chrysodeixis acuta</i>)</p>	<p>30 ml / 100 l water (3 ml / 10 l water)</p> <p style="text-align: center;">PLUS</p>	<p>Apply when the first larvae are observed – normally around the fruit-set stage of the crop. Apply at 500 to 1500 litres of spray mixture per hectare and ensure good coverage of all foliage. In the case of trellised tomatoes, apply to both sides of the tomato row. A suitable registered adjuvant such as Villa 51 can be added to improve efficacy. ADVANCE 150 SC will control larvae of all stages of development. Refer to “IMPORTANT NOTES” below.</p>
<p>Tomato leaf miner moth (<i>Tuta absoluta</i>)</p>	<p>50 ml / 100 l water (5 ml / 10 l water)</p> <p style="text-align: center;">Villa 51</p>	<p>Monitor the adult moth flights and commence with treatments at egg-hatching or when the very first erosions by the newly hatched larvae are visible. Apply at 500 to 1500 litres of spray mixture per hectare and ensure good coverage of all foliage. In the case of trellised tomatoes, apply to both sides of the tomato row. If required, apply follow up treatments at 7-day intervals up to a maximum of three (3) treatments per crop season. A suitable registered adjuvant such as Villa 51 can be added to improve efficacy.</p>
<p>IMPORTANT NOTES</p> <ul style="list-style-type: none"> • Regular scouting of tomato fields is important to establish the timing of the first treatment and subsequent applications if required. • ADVANCE 150 SC can be applied in an 8 to 14-day interval spray programme under conditions of continuous re-infestation of African bollworm, Tomato semi-looper or Potato tuber moth. • Use a 7-day spray interval for <i>Tuta absoluta</i>. • Do not exceed (3) treatments per season with ADVANCE 150 SC. The alternation of ADVANCE 150 SC with insecticides that have a different mode of action is recommended. • Apply a block application of two (2) consecutive ADVANCE 150 SC applications, and then switch to products with different modes of action. 		

Crop / Pest	Dosage rate	Remarks
<p>Wheat, Barley & Oats</p> <p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Ground Application:</u> 250 ml / ha</p> <p>OR</p> <p><u>Aerial Application:</u> 300 ml / ha</p> <p>PLUS</p> <p>a registered surfactant</p>	<p><u>Ground application:</u> Apply in 250 to 300 litres water per hectare in soybeans. Ensure thorough, even coverage of the foliage.</p> <p><u>Aerial application:</u> Apply in at least 30 litres water per hectare. Ensure thorough, even coverage of the foliage.</p> <p>Apply when the first bollworm eggs or small larvae (less than 10 mm in length) are observed following regular scouting. Larvae that are obscured by dense foliage during application may not be adequately controlled. It is therefore essential to ensure thorough coverage of the foliage and wheat ears. The addition of a suitable registered adjuvant, at the registered rate, is recommended to enhance efficacy. Do not apply more than one (1) application per season.</p>
<p>Tobacco</p> <p><u>Larvae of:</u> Potato tuber moth (<i>Phthorimaea operculella</i>)</p>	<p><u>Broadcast ground application:</u></p> <p>70 ml / 100 l water (175 to 210 ml /ha) (7 ml / 10 l water)</p> <p>ADVANCE 150 SC</p> <p>PLUS</p> <p>an organosilicon surfactant (such as Charge)</p>	<p>Apply after transplanting during the establishment phase of tobacco in the field as a preventative spray programme. Apply in 250 to 300 litres per hectare spray mixture at 7 to 10-day intervals with the first treatment 2 to 3 days after transplanting. Use the shorter interval under conditions of heavy infestation pressure. Direct the first and second applications as band applications, over the top of the plants, ensuring thorough coverage of the foliage, especially the lower third of the plants where Potato tuber moth larval infestations tend to be most severe. It is advised that the third and fourth application be directed from both sides. Use a boom fitted with drop arms with nozzles arranged to spray towards each other in order to improve coverage. Increase the spray volume as the plants grow in size. Apply a block application of 2 to 3 consecutive ADVANCE 150 SC applications, and then switch to products with other modes of action. Most beneficial insects and predatory mites are unaffected by applications of ADVANCE 150 SC. The addition of an organosilicon surfactant (such as Charge) is recommended to improve coverage.</p>
<p><u>Larvae of:</u> African bollworm (<i>Helicoverpa armigera</i>)</p>	<p><u>Ground application:</u></p> <p>250 ml / ha</p> <p>PLUS</p> <p>Villa 51</p>	<p><u>Field Treatment:</u> After transplanting into the field during the establishment and growing phases of tobacco. Apply ADVANCE 150 SC as a foliar application in 300 to 400 litres water per hectare. Increase the spray volume as plant size and density increases. Apply ADVANCE 150 SC early when bollworm eggs or small larvae are observed during regular scouting. Follow-up with a second ADVANCE 150 SC application at a spray interval of 10 to 14 days if necessary, under conditions of continuous re-infestation. Use the shorter interval under conditions of heavy infestation pressure. Do not exceed two (2) applications per season. The alternation of ADVANCE 150 SC with insecticides that have a different mode of action is recommended. Apply a block application of two (2) consecutive ADVANCE 150 SC applications, and then switch to products with different modes of action. Most beneficial insects and predatory mites are unaffected by applications of ADVANCE 150 SC. The addition of a non-ionic adjuvant (e.g., Villa 51) to the spray mixture is recommended to improve coverage.</p>

Crop / Pest	Dosage rate	Remarks
<u>Veld, Grazing, Cotton, Maize, Peppers & Soybeans</u> Fall army worm (<i>Spodoptera frugiperda</i>)	300 ml / ha PLUS 250 ml / ha Charge	Ground application: Apply in 200 to 400 litres water at the beginning of the infestation and at the first sign of damage. Ensure thorough coverage of the target area. Do not exceed two (2) consecutive applications of ADVANCE 150 SC per season. Use a spray interval of 7 days. Application is not recommended for control of caterpillars larger than 1 cm. The addition of a non-ionic surfactant (e.g., Villa 51) to the spray mixture is recommended to improve coverage.
	300 ml / ha	Aerial application: Apply in 30 litres water per hectare.
<u>Sorghum</u> Fall army worm (<i>Spodoptera frugiperda</i>)	300 to 400 ml / ha PLUS 250 ml / ha Charge	Apply as above.

Important note: Consult the **Lambda 50EC, Lambda Secure 106 CS, Villa 51, Aquabuff Plus, Interlock[®], Masterlock[®], Charge, Break-Thru[®] S240** and mineral oil adjuvant labels for **WARNINGS, PRECAUTIONS** and **DIRECTIONS FOR USE**.

The following products mentioned in this label may be replaced with equivalent products:

- **LAMBDA 50EC** (L 7787) = **JUDO 50 EC** (L 7785 / N-AR 1104) = **LAMBDA-CYTHRIN 50 EC** (L 8343) (**Lambda-cyhalothrin**),
- **LAMBDA SECURE 106 CS** (L 8939 / W 130693) = **JUDO SECURE 106 CS** (L 8938) = **MAGNUM SECURE 106 CS** (L 9172) (**Lambda-cyhalothrin**),
- **BARITONE 50 EC** (L 8659 / W 130692) = **PRIME 50 EC** (L 8660 / N-AR 1230) = **LUFRON 50 EC** (L 8916) (**Lufenuron**),
- **TEMPES 100 EC** (L 9946) = **RAMBA 100 EC** (L 9899) = **ZUMBA 100 EC** (L 9963) (**Bifenthrin**),
- **METHOMYL 90 SP** (L 5931 / W 130460) = **METHOMYL 90 SP** (L 7189 / N-AR 1084),
- **CHARGE** (L 9100) = **TECHNIWET SUPER** (L 9239) = **BREAK-THRU[®] S240** (L 6764),
- **VILLA 51** (L 8050 / W 130454 / N-AR 1090) = **WEN 51** (L 8315) and
- **AQUABUFF PLUS** (L 9210) = **TECHNIBUFF SUPER** (L 9209).

LAMBDA 50EC, BARITONE 50 EC, METHOMYL 90 SP, TEMPES 100 EC and/en **LAMBDA SECURE 106 CS** are registered products of / is geregistreerde produkte van **UNIVERSAL CROP PROTECTION (PTY) LTD.**

JUDO 50 EC, PRIME 50 EC, METHOMYL 90 SP, RAMBA 100 EC, JUDO SECURE 106 CS, TECHNIWET SUPER, VILLA 51, WEN 51, CHARGE, AQUABUFF PLUS and/en **TECHNIBUFF SUPER** are registered products of / is geregistreerde produkte van **VILLA CROP PROTECTION (PTY) LTD.**

LAMBDA-CYTHRIN 50 EC, LUFRON 50 EC, ZUMBA 100 EC and/en **MAGNUM SECURE 106 CS** are registered products of / is geregistreerde produkte van **CROP ASURE (PTY) LTD.**

BREAK-THRU[®] S240 is a registered product of / is 'n geregistreerde produk van **EVONIK DEGUSSA AFRICA (PTY) LTD.**

INTERLOCK[®] and/en **MASTERLOCK[®]** are registered products of / is geregistreerde produkte van **WINFIELD SOLUTIONS REGISTRATIONS HOLDINGS (PTY) LTD.**