

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Fungicide



CAPTAB WP

Reg. No. L 6758 Act/Wet No. 36 of/van 1947
N-AR 1903

1: 26/02/01-Oct2018

*A wettable powder fungicide for the control of diseases as listed.
'n Benatbare poeier swamdoder vir die beheer van siektes soos aangedui.*

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

captan 500 g/kg captan

FRAC FUNGICIDE GROUP CODE **M4** FRAC SWAMDODER GROEPKODE

kg

villa

Registration holder / Registrasiehouer:
Universal Crop Protection (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07
PO Box / Posbus 801, Kempton Park, 1620
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: 3077

Willow Set & Print 011 394-4486



**CAUTION
VERSIGTIG**



GEBRUIKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED
REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

CAPTAB WP

Reg. No. L 6758 / N-AR 1903 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: M4

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

captan / kaptan 500 g/kg

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620

Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS****Withholding periods:**

| Allow the following withholding period between the last application and harvest or feeding: | |
|--|--------------|
| Celery | 3 days |
| Tomatoes | 7 days |
| Strawberries | 10 days |
| Apples, Apricots, Boysenberries, Youngberries, Guavas, Olives, Peaches, Pears, Plums and Quinces | 14 days |
| Table grapes (do not apply after pea size) | 4 to 6 weeks |

IN THE CASE OF CROPS INTENDED FOR EXPORT, CONFIRM WITHHOLDING PERIODS WITH THE RELEVANT EXPORT ORGANISATION. IF THIS PRODUCT IS USED ACCORDING TO THE RECOMMENDATIONS ON THIS LABEL, LOCAL MRL VALUES WILL NOT BE EXCEEDED. HOWEVER, UNIVERSAL CROP PROTECTION (PTY) LTD. CANNOT ACCEPT RESPONSIBILITY FOR EXPORTED CROPS EXCEEDING THE IMPORT TOLERANCES OF OTHER COUNTRIES.

- Handle with care.
- May cause eye irritation and skin sensitization.
- Harmful if inhaled or swallowed.
- Toxic to fish and wildlife.
- Store away from food and feeds.
- Store in the original container/bag, tightly closed, in a cool, dry place.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the diseases against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Do not inhale the dust or spray mist.
- Avoid contact with skin, eyes and clothing.
- Wash with soap and water after use and accidental skin contact.
- Do not eat, drink or smoke whilst applying, mixing, before washing hands and face or change of clothing.
- Prevent spray drift onto crops, grazing, rivers, dams or areas not under treatment or adjacent water sources.
- Clean the applicator after use and dispose of wash water where it will not contaminate food, crops, grazing, rivers or dams.
- Containers and packages must be completely emptied before being disposed of.
- Shake out thoroughly into the mixing or spray tank before destroying the empty container/bag.

- **Never** re-use the container/bag for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

- **CAPTAB WP** is compatible with and may be sprayed together with most commonly used insecticides, fungicides and trace elements.
- Not compatible with alkaline materials such as Lime sulphur.
- It is however recommended to perform physical compatibility and crop safety tests on a limited scale, before a tank mixture is sprayed.

Mixing instructions:

- Make sure the spraying equipment is clean, before use.
- Half fill the spray tank and start the agitator.
- Mix the required quantity of **CAPTAB WP** with a little water into a thin paste and add to the water in the spray tank, while stirring.
- Stir continuously, to ensure a homogenous mix and add the rest of the required volume of water.
- Keep agitating whilst applying.

Application instructions:

- Unless otherwise stated, spray **CAPTAB WP** at 7 to 10 day intervals and after rain.
- Apply to the point before runoff.

APPLICATION RATES

| <i>Crop / Disease</i> | <i>Dosage rate</i> | <i>Remarks</i> |
|--|---------------------|--|
| Apples and Pears Post harvest decay | 200 g / 100 ℓ water | Apply as a wash to walls and floors to aid disinfection. Only for use on empty premises. |
| Scab | 200 g / 100 ℓ water | Do not use in combination with Sulphur or after a Lime sulphur application, especially on White Winter Pearmain. Pre-blossom: Apply the first application at green tip stage and repeat at 7 day intervals. Post-blossom: Continue with sprays. Apply 3 to 4 sprays according to the weather conditions, at 10 to 14 day intervals. |
| NOTE CAPTAB WP may cause Leaf spotting of Pearmain, Delicious, Cox and other sensitive apple varieties. | | |
| Apricots and Peaches Freckle | 200 g / 100 ℓ water | Winter rainfall region: Apply the first application at 75 to 100 % petal drop. Thereafter repeat the application 14 days later. During wet seasons, repeat the application monthly on late cultivars. Summer rainfall region: Apply the first application 3 weeks after petal drop and apply the second application 14 days later. During wet seasons repeat the application monthly on late cultivars. |

| Crop / Disease | Dosage rate | Remarks |
|---|--|---|
| <u>Apricots and Plums</u> Gum spot | 200 g / 100 ℓ water | Spray the first application at 95 to 100 % petal drop and repeat 14 days later. Green rot on apricots will also be controlled. |
| <u>Peaches and Plums</u> Bacterial spot | 200 g / 100 ℓ water | The level of infection of the previous season will determine if one or two applications at 50 % and/or 75 % leaf drop would be required. If twig and bud infections are observed before bud movement, apply an additional spray at bud movement. Persist with post blossom applications, at 10 to 14 day intervals, up to 3 weeks before picking. |
| <u>Boysenberries and Youngberries</u> Downy mildew | 100 g / 100 ℓ water | Apply 4 sprays at 10 to 14 day intervals. Start with the first application 14 days after green-tip stage. |
| <u>Celery</u> Leaf spot | 200 g / 100 ℓ water | Apply as a full cover spray. When the infestation is severe, repeat the application after 10 days. |
| <u>Guavas</u> Blossom-end rot | 200 g / 100 ℓ water | Apply when very wet weather conditions are experienced, for protection against Blossom-end rot. Apply as a full cover spray 14 days before harvest. |
| <u>Lawns</u> Dollar spot, Fairy ring, Crown & Root rot diseases | 300 g / 100 ℓ water | Apply at the first signs of diseases. Apply 200 litres spray mixture per 100 m ² . |
| <u>Olives</u> Leaf spot | 200 g / 100 ℓ water | Apply 2 applications. Apply the first spray in spring, before the new flush appears and the second spray in autumn, before the start of winter. |
| <u>Potatoes</u> Late blight | 200 to 300 g / 100 ℓ water (2 to 3 kg / ha) | Start spraying before the disease is due to appear or when symptoms of the disease appear. Apply every 7 to 10 days if weather conditions are favourable for disease development. Use the higher dosage rate for big plants and/or when the infection is severe. The dosage rate depends on the severity of the infection. Apply 500 to 1500 litres spray mixture per hectare, depending on the plant size and density. For Low Volume application, adjust the spray tank concentration, to apply not more than 3 kg per hectare. Ensure thorough coverage of the upper and lower leaf surfaces. Repeat the application after more than 13 mm of rain or overhead irrigation. |
| <u>Quinces</u> Black spot | 200 g / 100 ℓ water | Commence the applications at green point stage and repeat every 6 to 10 days. Lengthen the spray interval between applications from 10 to 14 days at 75 % petal drop stage. |
| <u>Roses</u> Black spot | 200 g / 100 ℓ water | Apply as a full cover spray. Commence the applications when the first symptoms of the disease are noticed. Repeat the applications at weekly intervals. Do not spray when dry weather conditions are experienced. |
| <u>Strawberries</u> <i>Botrytis</i> rot | 100 g / 100 ℓ water | Commence the applications at 5 % flowering stage, when the disease is noticed. Repeat the applications every 7 to 10 days. |
| Leaf spot | 100 g / 100 ℓ water | Apply every 10 to 14 days, depending on the severity of the disease. |

| Crop / Disease | Dosage rate | Remarks |
|--|---|---|
| <u>Table Grapes</u> Anthracnose | 200 g / 100 ℓ water (0.5 to 1 kg / ha) | Do not apply after berries have reached pea-size (4 to 6 weeks prior to harvest), as this may cause tarnishing. Apply 250 to 500 litres spray mixture per hectare. Summer rainfall region: If rains commence September, apply the first application shortly before flowers open. Alternatively apply to young re-growth and bunches every 14 days, after flowering, until berries start swelling and become translucent. Winter rainfall region: Apply the first application when shoots are 1 to 2 cm long and repeat when shoots are 10 cm long. Thereafter apply a third and fourth application at 2 to 3 week intervals. |
| <u>Tomatoes</u> Late blight & Septoria leaf spot | 200 g / 100 ℓ water (2 to 3 kg / ha) | Seedbeds: Spray twice a week. Lands: Start the application shortly after transplant, when plants are about 15 cm high and before the disease is due to appear. Apply 500 to 2000 litres spray mixture per hectare, depending on the plant size and density. Apply every 7 to 10 days if the weather conditions are favourable for disease development. Use the higher dosage rate for big plants and/or when the infection is severe. The dosage rate depends on the severity of the infection. For Low Volume application, adjust the spray tank concentration to apply not more than 3 kg per hectare. Ensure thorough coverage of the upper and lower leaf surfaces. Repeat the application after more than 13 mm rain or overhead irrigation. |
| <u>Tobacco</u> Seedbed diseases (Anthracnose, Black root rot, Black shank & Damping off) | 200 g / 100 ℓ water | Apply 15 litres mixture per 10 m ² as a drench treatment, 3 weeks after germination. Thereafter repeat the treatment every 7 to 14 days. After treatment apply 40 litres water to the treated area, to wash the product into the soil. |
| <u>Soils and Potting mixtures</u> Damping off, Seed decay & Root rot | 400 g / 100 ℓ water | Apply 20 litres spray mixture per 10 m ² to the seedbed, prior to sowing. Thoroughly incorporate the spray mixture with the top 5 to 10 cm of the soil layer. Alternatively spray 5 litres spray mixture per 10 m ² directly onto the soil, after sowing. |