

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Fungicide



**COPPASULFA**

Reg. No. L 5726 Act/Wet No. 36 of/van 1947

1:27/6/2017 – June 2017

*A dusting powder for the control of diseases and mites on crops as listed.  
'n Stuifpoeier vir die beheer van siektes en myte op gewasse soos gelys.*

**ACTIVE INGREDIENTS / AKTIEWE BESTANDEDELE**

copper oxychloride	<b>85 g/kg</b>	koperoksiechloried
sulphur	<b>900 g/kg</b>	swael

FRAC FUNGICIDE GROUP CODE **M 1/2** FRAC SWAMDODER GROEPKODE

kg

**villa**

Registration holder / Registrasiehouer:

Villa Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1992/002474/07

PO Box / Posbus 10413, Aston Manor, 1630

Tel: 011 396 2233

Website / Webblad: [www.villacrop.co.za](http://www.villacrop.co.za)

UN Number: 1350

Willow Set & Print 011 394-4486



CAUTION  
VERSIGTIG



GEBRUIKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE  
GEDRUK OP HOUER/SAK

Date formulated:  
Formuleringsdatum:

REFER TO DETAILS PRINTED  
ON CONTAINER/BAG

DIRECTIONS FOR USE ENCLOSED

Batch number:  
Lotnommer:

**COPPASULFA**

Reg. No. L 5726 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: M 1/2

**ACTIVE INGREDIENT / AKTIEWE BESTANDELE:**

copper oxychloride / koperoksiechloried ..... 85 g/kg  
 sulphur / swael .....900 g/kg

Registration holder / Registrasiehouer:

**VILLA CROP PROTECTION (PTY) LTD.**

Co. Reg. No. 1992/002474/07 Mpy. Reg. Nr.

PO Box / Posbus 10413

ASTON MANOR, 1630

Tel. (011) 396 2233

**CAUTION / VERSIGTIG****WARNINGS**

Allow the following waiting periods between last application and harvests:	
Cucurbits, Table Grapes and Tomatoes	3 days
Wine Grapes	4 to 6 weeks

**IN THE CASE OF CROPS INTENDED FOR EXPORT, CONFIRM WITHHOLDING PERIODS WITH THE RELEVANT EXPORT ORGANISATION. IF THIS PRODUCT IS USED ACCORDING TO THE RECOMMENDATIONS ON THIS LABEL, LOCAL MRL VALUES WILL NOT BE EXCEEDED. HOWEVER, VILLA CROP PROTECTION (PTY) LTD. CANNOT ACCEPT RESPONSIBILITY FOR EXPORTED CROPS EXCEEDING THE IMPORT TOLERANCES OF OTHER COUNTRIES.**

- Keep out of reach of children, uninformed persons and animals.
- Toxic to fish.
- Store in a cool, dry place, away from food and feeds.
- **Flammable – keep away from flames and sparks.**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease/mites to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

**PRECAUTIONS**

- Do not inhale dust.
- Do not eat, drink or smoke whilst applying.
- Prevent contamination of food, feeds, drinking water and eating utensils.
- Avoid drift to other crops, grazing, rivers and dams.
- Destroy empty container in such a way that it cannot be used for any other purpose.

**USE RESTRICTIONS**

- Heavy applications may cause "burn" on crops, especially on hot days.
- When table grapes have reached pea-size stage, do not apply while bunches are still wet with free water following e.g. rain or heavy dew.

**DIRECTIONS FOR USE: Use only as directed.**

Dust plants lightly by means of a suitable applicator. Repeat after rain. Dosage depends on plant size.

**APPLICATION RATES**

**Important - refer to "USE RESTRICTIONS" above.**

<b>Crop / Disease</b>	<b>Dosage</b>	<b>Remarks</b>
<b>Cucurbits</b> Downy mildew & Powdery mildew	10 to 30 kg	<b>Do not use on muskmelon or cucumber.</b> Start application at first sign of disease. Repeat weekly as long as weather favors the disease. Ensure coverage of both leaf surfaces.
<b>Grapes</b> Anthracnose, Bacterial blight, Downy mildew, Powdery mildew, Erinose mite (bud mite) and Dead arm	15 to 30 kg	First application when shoots are 1 to 2 cm long and again when shoots are 10 cm long. Thereafter apply 2 to 3 times at 2 to 3 week intervals.
Botrytis rot (mainly on table grapes)	15 to 30 kg	Apply 6 weeks before bunches ripen and repeat at 7 to 10 day intervals. Ensure a light even coverage of bunches.
<b>Mangoes</b> Powdery mildew	20 to 40 kg	Apply at first sign of infection and at full blossom. Repeat if blossoming is irregular and after heavy rain.
<b>Ornamentals &amp; Flowers</b> Powdery mildew & Downy mildew	10 to 30 kg	Commence dusting at the first sign of the disease and repeat every 7 to 14 days depending on weather conditions. Ensure complete coverage.
<b>Roses</b> Powdery mildew & Black spot	15 to 30 kg	Apply at first sign of disease and repeat weekly. Application must be discontinued when a dry period is experienced.
<b>Tomatoes</b> Early blight & Septoria leafspot	15 to 30 kg	Commence dusting shortly after transplant when plants are about 15 cm high and before symptoms appear. Repeat every 4 to 10 days depending on weather conditions. Will also control Rust mite and Erinose mite.