

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE

Insecticide



DICLOFEN 240 SC

Reg. No. L 9497 Act/Wet No. 36 of/van 1947

1:11/3/2014-Feb2017

A contact suspension concentrate acaricide for the control of various mite pests on citrus.

'n Kontak suspensie konsentraat mytdoder vir die beheer van verskeie mytplae op sitrus.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

spirodiclofen (tetronic acid) **240 g/l** spirodiklofen (tetroniese suur)

IRAC INSECTICIDE GROUP CODE **23** IRAC INSEKODODER GROEPKODE



villa

Registration holder / Registrasiehouer:

Universal Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07

PO Box / Posbus 801, Kempton Park, 1620

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: 3082

Willow Set & Print 011 394-4486



CAUTION
VERSIGTIG



GEBRUIKSAANWYINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

DIRECTIONS FOR USE ENCLOSED

DICLOFEN 240 SC

Reg. No. L 9497 Act/Wet No. 36 of/van 1947

IRAC INSECTICIDE GROUP CODE / IRAC INSEKODER GROEPKODE: 23

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

spirodiclofen (tetronic acid)

.....240 g/l

spirodiklofen (tetroniese suur)

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801

KEMPTON PARK, 1620 Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS****Withholding periods:**

Allow the following number of days between last application and harvest of the following crops:	
Citrus	76 days

IN THE CASE OF CROPS INTENDED FOR EXPORT, CONFIRM WITHHOLDING PERIODS WITH THE RELEVANT EXPORT ORGANISATION. IF THIS PRODUCT IS USED ACCORDING TO THE RECOMMENDATIONS ON THIS LABEL, LOCAL MRL VALUES WILL NOT BE EXCEEDED. HOWEVER, UNIVERSAL CROP PROTECTION (PTY) LTD. CANNOT ACCEPT RESPONSIBILITY FOR EXPORTED CROPS EXCEEDING THE IMPORT TOLERANCES OF OTHER COUNTRIES.

- Handle with care.
- Harmful if swallowed, inhaled or absorbed through skin. Prevent eye contamination.
- Toxic to fish and aquatic invertebrates.
- Store under lock and key, in a cool, dry, well-ventilated place.
- Store away from food, feeds, seed, fertilizers and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.
- **In case of poisoning, take the patient immediately to a doctor and make this label available to him/her.**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions. The action and effect thereof may be effected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Avoid contamination to surface water through spray drift. Do not contaminate water when disposing of equipment wash water or rinsate. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

This product is toxic to honeybee larvae through direct contamination of pollen and nectar. Do not apply to blooming, pollen-shedding or nectar-producing parts of plants if bees forage on the plants.

SURFACE WATER ADVISORY AND RUN-OFF MANAGEMENT

This product may contaminate water through run-off or drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce run-off that contains this product. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall run-off.

Run-off of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

PRECAUTIONS

- Avoid inhalation of the spray mist or fumes.
- Avoid eye and skin contact.
- Wear a face shield, rubber gloves and boots, when handling the concentrate and preparing the spray mixture.
- Wash with soap and water after use and after accidental skin contact.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face and change of clothing.
- Prevent drift on to other crops, grazing, rivers, dams or areas not under treatment or to nearby water sources.
- Thoroughly clean spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank, before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening and dispose of it in a safe way.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

RESISTANCE WARNING

DICLOFEN 240 SC is a group code 23 acaricide/insecticide. Any mite population may contain individuals naturally resistant to **DICLOFEN 240 SC** and other group code 23 acaricides/insecticides. The resistant individuals can eventually dominate the mite population if these acaricides/insecticides are used repeatedly. These resistant mites may not be controlled by **DICLOFEN 240 SC** or any other group code 23 acaricide/insecticide.

To delay acaricide resistance:

- avoid exclusive repeated use of acaricide from the same acaricides/insecticides group code. Alternate or tank mix with products from different acaricides/insecticides group codes,
- integrate other control methods (chemical, cultural, biological) into mite control programmes.

For specific information on resistance management contact the registration holder of this product.

USE RESTRICTIONS:

- Do not use brackish water with **DICLOFEN 240 SC**.
- Only apply 1 treatment per season.
- **Citrus:** Do not apply **DICLOFEN 240 SC** later than mid-January and within 76 days of harvest.

GENERAL INFORMATION

- **DICLOFEN 240 SC** is a non-systemic acaricide and is active by contact on mite eggs, all nymphal stages and adult females. Apply **DICLOFEN 240 SC** when mite populations are building up or timed to coincide with early threshold level in developing mite populations and not as a corrective treatment. Thorough coverage of the leaf upper and lower surfaces and of the fruit is therefore essential, to ensure long lasting efficacy. The presence of physical barriers including, but not limited to excessive dust, sun block agents, and/or sooty-mold can interfere with the binding of **DICLOFEN 240 SC** to target plant parts and may require higher spray volumes to achieve penetration of these barriers.

Restrictions

- Do not apply **DICLOFEN 240 SC** through any type of irrigation system;
- Do not apply **DICLOFEN 240 SC** in enclosed structures such as greenhouses or plant houses.
- **Effect on beneficial insects in citrus**
Aphytis lingnanensis - Harmless
Chilocorus nigritus – Slightly harmful
Coccidoxenoides peregrinus – Harmful / Slightly harmful
Euseius citri - Harmless

Trichogrammatoidea cryptophlebiae - Harmless
Bees - Harmless

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

- The formulation of other products as well as the quality of the dilution water may influence the compatibility with **DICLOFEN 240 SC**. A physical compatibility test should thus always be carried out before spraying. The Registration Holder cannot accept responsibility for any adverse effects due to tank mixtures with any product(s) not approved in this label.
- **DICLOFEN 240 SC** is **NOT** compatible with fungicides containing **Trifloxystrobin**.

Mixing Instructions:

- Half fill the spray tank with clean water and ensure agitation.
- Add the required volume of **DICLOFEN 240 SC** to the water.
- Agitate the water in the spray tank and then add the product(s) to the tank in the following sequence (as applicable): acidifier/buffer or adjuvant, suspension concentrate, water soluble concentrate, emulsifiable concentrate.
- Fill the spray tank with water to the required level while maintaining agitation, to ensure thorough mixing.
- Maintain agitation while spraying.
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

Application Instructions:

Ground application:

- It is important that all applications must be performed with suitable equipment that is in good working order and correctly calibrated, to give the desired coverage for that particular method of application.
- It is important to ensure complete coverage of the fruit and foliage to obtain maximum long lasting efficacy.
- Application rates specified within this label are based on full-size mature trees.

APPLICATION RATES

<i>Crop / Pest</i>	<i>Dosage Rate</i>	<i>Remarks</i>
Citrus Excluding lemons and kumquat Citrus red mite, <i>(Panonychus citri)</i> Citrus flat mite, <i>(Brevipalpus californicus)</i> Lowveld citrus mite, <i>(Eutetranychus orientalis)</i>	10 to 15 ml / 100 l water	Apply as a medium cover spray when first signs of an infestation are observed. Only apply 1 treatment per season before middle of January. Precede, and then follow the DICLOFEN 240 SC treatment with registered acaricides with other modes of action, if necessary. NOTE Do not apply Trifloxystrobin 500 WDG within 30 days before or after the application of DICLOFEN 240 SC .
Citrus rust mite, <i>(Phyllocoptruta oleivoar)</i>	15 to 20 ml / 100 l water	

IMPORTANT NOTES

Apply of **DICLOFEN 240 SC** as a spray early in the infestation as the Mite populations begin to develop or at an early threshold level for the target mite species. Evaluate the product's performance no sooner than 7 days following application. Take note that only mite eggs, all nymphal stages and adult females will be controlled by the product – **NOT** the adult males. The degree of control or suppression of additional labelled pests will be determined, in part, by the stage of pest development at application and infestation level of those pests.

Use higher rates for moderate to heavy Mite pressure or where longer residual control is desired. Lower rates are generally adequate for low to moderate Mite pressure, but require careful scouting and will generally provide shorter residual control than higher rates.

Apply adequate spray volume to ensure thorough and uniform coverage of target plant parts.