

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Insecticide



ELIREX®

Reg. No. L 8657 Act/Wet No. 36 of/van 1947

3/26/11/2013-June2017

*A bait in pellet form for control of Snails in agricultural crops and horticulture.*

*'n Lokaas in korrelvorm vir die beheer van Slakke in landbougewasse en tuinbou.*

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

metaldehyde 50 g/kg metaldehid

20kg

villa 

Registration holder / Registrasiehouer:  
Universal Crop Protection (Pty) Ltd.  
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07  
PO Box / Posbus 801, Kempton Park, 1620  
Tel: 011 396 2233  
Website / Webblad: [www.villacrop.co.za](http://www.villacrop.co.za)

UN Number: Not Regulated

Willow Set & Print 011 394-4486



CAUTION  
VERSIGTIG



GEBRUIKSAANWYSINGS INGESLUIT

VERWYS NA BESONDERHEDE  
Date formulated: GEDRUK OP HOUER/SAK  
Formuleringsdatum: GEDRUK OP HOUER/SAK

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED  
ON CONTAINER/BAG  
Batch number:  
Lotnummer:

**ELIREX®**

Reg. No. L 8657 Act/Wet No. 36 of/van 1947

**ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:**

metaldehyde / metaldehyd ..... 50 g/kg

Registration holder / Registrasiehouer:

**UNIVERSAL CROP PROTECTION (PTY) LTD.**

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

PO Box / Posbus 801

KEMPTON PARK, 1620.

Tel. (011) 396 2233

**CAUTION / VERSIGTIG****WARNINGS**

- **Withholding periods:**
  - Do not harvest edible crops within 7 days after application.
  - Do not allow livestock or poultry to graze treated areas.
- Handle with care.
- Harmful if swallowed or inhaled.
- Toxic to fish and wildlife.
- Store the product inside the original container/bag and always keep the container/bag tightly closed.
- Store in a cool, dry, well-ventilated place, away from food, feeds, seed and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.

**Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.**

**PRECAUTIONS**

- Avoid contact with the skin and eyes.
- Wear rubber gloves when applying the product.
- Do not eat, drink or smoke whilst preparing or applying or before washing hands and face or change of clothing.
- Do not apply near water bodies such as fishponds.
- Container/bags must be completely emptied before being disposed of. Shake out thoroughly and destroy the empty container/bag thereafter and bury it. **Do not burn the empty container/bag.**
- **Never** re-use the empty container/bag for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

**DIRECTIONS FOR USE: Use only as directed.****Use restrictions:**

- Do not apply to edible plant parts or crops where baits may be collected with harvested commodities.
- Do not heap baits.
- To prevent accidental, but fatal poisoning, product should be kept away from ducks, poultry, animals and domestic pets. Contact a veterinary surgeon if pets or livestock are poisoned.

**General information:**

- **ELIREX®** kills Snails by irreversibly destroying the mucus production cells that are essential for movement.
- Under high infestation pressure it may be necessary to repeat application.
- Best results are achieved by application during mild damp weather when Snails are most active, but when rain is not imminent.
- For seedlings, apply 5 to 10 days prior to planting.

**APPLICATION RATES**

| <b>Situation</b>  | <b>Application rate</b>   | <b>Remarks</b>  |
|---|---|---|
| Apply bait when crops are most vulnerable and when the pest is evident and active. Generally start when rain/moisture triggers Snail activity, but before significant egg laying occur. |   |   |
| <b><u>Crops, Orchards and Vines</u></b><br>Brown snail<br>( <i>Helix aspersa</i> )  | 5 to 8 kg / ha  | Apply as a broadcast application evenly over the ground surface where Snails are active. In grapevines, also apply close to supportive struts.<br>Repeat application if infestation re-appears.     |
| <b><u>Vines</u></b><br>Dune snail<br>( <i>Theba pisana</i> )  | 5 to 7.5 kg / ha  | Apply from budding stage as a broadcast application evenly over the ground surface where Snails are active. Also apply close to supportive struts.<br>Repeat application if infestation re-appears. |
| <b><u>Horticulture &amp; Gardens</u></b><br>Brown snail<br>( <i>Helix aspersa</i> )   | 5 to 8 g per 10 m <sup>2</sup> area<br><br><b>OR</b><br>35 to 40 pellets per m <sup>2</sup> | Scatter the <b>ELIREX</b> <sup>®</sup> bait pellets evenly in flower beds, under dense foliage, shrubs and trees. In case of container plants, place bait pellets close to base of container.       |

**ELIREX**<sup>®</sup> is a registered trade mark of / is 'n geregistreerde handelsmerk van  
**DE SANGOSSE**