



A WATER-SOLUBLE CONCENTRATE PLANT GROWTH REGULATOR
FOR FRUIT THINNING ON APPLES.

2: 6/9/2011 – June 2017

'N WATEROPLOSBAARE KONSENTRAAT PLANTGROEIREGULEERDER
VIR UITDUN VAN VRUGTE OP APPELS.

REG. NO. L 7891 ACT/WET NO. 36 OF/VAN 1947

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

6-BENZYLAMINOPURINE (CYTOKININ) / 6-BENSIELAMINOPURIEN (SITOKINIEN)21 G/l

UN NUMBER: NOT REGULATED

MANUFACTURED BY /
VERVAARDIG DEUR: **FINE AGROCHEMICALS LIMITED**
HILL END HOUSE WHITTINGTON
WORCESTER WR5 2RQ UK
ENQUIRE@FINE.EU
WWW.FINE.EU

DISTRIBUTED BY /
VERSPREI DEUR:



REGISTRATION HOLDER/REGISTRASIEHOUER
UNIVERSAL CROP PROTECTION (PTY) LTD.
CO. REG. NO. 1983/008184/07
MPY. REG. NR. P.O.BOX / POSBUS 801
KEMPTON PARK, 1620
TEL. (011) 396 2233
WEBSITE / WEBBLAD: WWW.VILLACROP.CO.ZA

BATCH NUMBER /
LOTNOMMER:

DATE FORMULATED /
FORMULERINGSDATUM:

NET VOLUME:
NETTO VOLUME:

5l

EX/SA/5L/17

fine

© EXILIS® IS A TRADEMARK OF / IS 'N HANDELSMERK VAN FINE AGROCHEMICALS LTD., UK.

**CAUTION
VERSIGTIG**



Vir volledige besonderhede, sien ingeslote pamflet.
For full particulars, see enclosed leaflet.

EXILIS®

Reg. No. L 7891 Act/Wet No. 36 of/van 1947

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

6-benzylaminopurine (cytokinin) / 6-bensielaminopurien (sitokinien)..... 21 g/l

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O.Box / Posbus 801,

KEMPTON PARK, 1620

Tel. (011) 396 2233

WARNINGS

- Harmful if swallowed.
- Store in a cool place, away from food, feeds, seed, fertilizers and other agricultural remedies.
- Can cause eye irritation.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Do not inhale the spray mist.
- Avoid contact with skin, eyes and clothing.
- Wash with soap and water after use.
- In case of contact with eyes, flush with plenty of water and get medical attention.
- In case of contact with skin, wash immediately with soap and plenty of water and remove contaminated clothes.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face or change of clothing.
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean the applicator after use and dispose of wash water where it will not contaminate food, grazing, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the container by perforation and flattening and dispose of it in a safe way.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

DIRECTIONS FOR USE: Use only as directed.**Compatibility:**

- Do not apply **EXILIS®** in combination with pesticides and other growth regulators.
- It is advisable to add a buffer e.g. **COMMODOBUFF** (L 5390 / N-AR 1107) to alkaline spray water to reduce the pH below 7.

Application Instructions:

- **EXILIS®** is a plant growth regulator for use on apples as a post-bloom thinning agent.
- **EXILIS®** must be absorbed by the plant to be effective.

- Optimum results will be achieved on vigorous, healthy trees when sprayed under good absorption conditions.
- High relative humidity and slow drying conditions favour maximum absorption.
- Do not apply **EXILIS®** to low vigor trees, or trees under stress from drought, winter injury etc.
- Apply as a uniform fine mist spray so that 85 % of the spray volume is directed into the upper two thirds of the trees under calm weather conditions.
- Use a properly calibrated sprayer that ensures uniform and complete coverage of all foliage and blossoms. Spray volume should be adjusted to tree size and density. Spray volumes around 1000 litres per hectare should be adequate.
- Do not apply **EXILIS®** when the air temperature is lower than 10 °C or greater than 32 °C. Rapid drying conditions should be avoided. Applications in the mornings or evenings may be preferable.
- Temperatures of over **18°C** for the two days following application are considered important for good thinning. High temperatures (+28 °C over this period) may result in excessive thinning.
- Delay **EXILIS®** application after rain until flower parts and leaves are free of surface droplets. Rainfall within 6 hours of application may reduce the expected response of **EXILIS®**.
- Thinning responses are normally to be observed within 10 to 14 days of application.

Mixing Instructions:

- Add the required amount of **EXILIS®** to a **clean** spray tank half filled with water.
- Agitate while filling the tank with water to the required level.
- **EXILIS®** spray solutions should be used within 24 hours of mixing.
- When a buffered wetting agent is used, it should be added to the tank prior to the addition of **EXILIS®**.

APPLICATION DOSAGE RATES

<i>Crop</i>	<i>Growth Regulating Activity</i>	<i>Dosage Rate ml / 100 l water</i>	<i>Remarks</i>
Apples (Golden Delicious, Royal Gala, Pink Lady, Granny Smith)	Fruit thinning	500 to 750 ml (in a minimum spray volume of 1000 to 1200 l / ha)	Apply EXILIS® when the king fruitlets are between 10 to 12 mm, but not later than 30 days after full bloom.

EXILIS® is a trademark of / is 'n handelsmerk van
FINE AGROCHEMICALS LTD., UK.

COMMODOBUFF is a registered product of / is 'n geregistreerde produk van
VILLA CROP PROTECTION (PTY) LTD.

PRODUCT WARRANTY CONDITIONS AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications indicated on this label. FINE disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance or other unintended consequences may result because of factors such as abnormal weather conditions, presence of other materials, the manner of application, use of the Product other than in strict accordance with this label's instructions or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, the Buyer shall assume all such risks.

FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product. FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product.

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ, United Kingdom.