



falgro[®] TABLETS

5:15/07/04 - June 2017

A PLANT GROWTH HORMONE IN TABLET FORM FOR THE PROMOTION OF CERTAIN PHYSIOLOGICAL AND MORPHOLOGICAL EFFECTS AS INDICATED ON SPECIFIC CROPS.

'N PLANT-GROEIHORMOON IN TABLETVORM VIR DIE BEVORDERING VAN SEKERE FISILOGIESE EN MORFOLOGIESE EFFEKTE SOOS AANGEDUI OP SPESIFIEKE GEWASSE.

REG. NO. L 4503 ACT/WET NO. 36 OF/VAN 1947

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

GIBBERELIC ACID / GIBBERELLIENSUUR 204 G/KG (1 TABLET = 1 G A.I. = 1 PPM)

4.9 g x 10 Tablets

BATCH NUMBER:

LOTNUMMER:

DATE FORMULATED:

FORMULERINGSDATUM:

UN NUMBER: NOT REGULATED

Manufactured by/
vervaardig deur:

fine

FINE AGROCHEMICALS LIMITED

Hill End House Whittington
Worcester WR5 2RQ UK
E enquire@fine.eu
www.fine.eu

Distributed by /
Versprei deur:

villa

Registration holder / Registrasiehouer:
UNIVERSAL CROP PROTECTION (PTY) LTD.
Co. Reg. No. 1983/008184/07 Mpy. Reg.
KEMPTON PARK, 1620
Tel. (011) 396 2233
Website/Webblad: www.villacrop.co.za

© FALGRO IS A REGISTERED TRADEMARK OF / IS 'N GEREJISTREERDE HANDELSMERK VAN FINE AGROCHEMICALS LTD.



CAUTION / VERSIGTIG



Vir volledige besonderhede, sien ingeslote pamflet. - For full particulars, see enclosed leaflet.

FAJTA/ZA/17

FALGRO® TABLETS

Reg. No. L 4503 Act/Wet No. 36 of/van 1947

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

gibberellic acid / gibberelliensuur 204 g/kg (1 tablet = 1 g a.i. = 1 ppm)

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620, Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS**

- Handle the product with care.
- May cause mild skin irritation and slight eye irritation.
- Harmful when swallowed.
- Store in a cool, dry, well-ventilated place.
- Store away from food, feeds, seed, fertilizers and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Do not inhale the spray mist.
- Avoid contact with skin and eyes.
- Wash with soap and water after use and after accidental skin contact.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face and change of clothing.
- Do not allow drift of spray onto other crops, grazing, rivers, dams or areas not under treatment.
- Packaging and containers must be completely emptied before being disposed of. Shake out thoroughly into the mixing tank and destroy the empty packaging in a safe way. **Never** re-use the empty packaging and containers for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.
- Clean the applicator and dipping bath after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.

USE RESTRICTIONS

- The application of gibberellic acid to certain **citrus** cultivars, such as Turkey Valencia and Bhihinnia Navel, may cause phytotoxicity.
- Do not apply **2,4-D Ester** in **citrus** destined for export before consulting the export guidelines.
- Do not use **FALGRO™ TABLETS** on **Sultanas**, which are to be dried.
- Do not apply to **vines**, which are suffering from drought.

DIRECTIONS FOR USE: Use only as directed.

- Gibberellic acid performs most effectively when the spray/dip solution pH is 5.5 to 6.5. Higher pH values may impair product performance, whilst lower pH values may affect the crop negatively. It is recommended that the spray/dip solution be buffered, if the pH is outside 5.5 to 6.5.
- Environmental factors influence the efficacy of gibberellic acid.
- Water stress and heat waves can have negative influences.
- It is recommended that soil water conditions should be at field capacity during application.

- Applications should be early in the morning or late in the afternoon. Do not apply during the heat of the day.

Compatibility:

- Compatible with most normal spray materials, except for mineral oils and alkaline mixtures like Bordeaux mixture and Lime Sulphur.
- Do not use chlorinated water.
- Do not apply with other agricultural chemicals not indicated on this label.
- Citrus: If a summer oil has been sprayed on citrus, a minimum of 14 days must elapse before **FALGRO® TABLETS** may be applied. Oil may be applied 5 to 10 days after a **FALGRO® TABLETS** application.

Mixing instructions:

- Dissolve the required quantity of sachets in a small quantity of water.
- Ensure sachets are completely dissolved before adding the solution to the spray tank or dipping bath.
- Fill the spray tank/dipping bath with water to the required level, while maintaining agitation to ensure thorough mixing.
- Ensure the pH of the spray/dip solution is between 5.5 to 6.5. If not, add the suitable buffering agent to the solution before using.
- Maintain agitation during application.
- **IMPORTANT NOTE**
FALGRO® TABLETS are not very stable in water and spray mixtures should definitely not be left overnight.

APPLICATION RATES

<i>Crop / Use</i>	<i>Dosage Rate</i>	<i>Remarks</i>
<u>Grapes</u>		Sultana Grapes: Apply in 1300 to 1800 litres water per hectare. Ensure thorough wetting of bunches and enclosing leaves. Care must be taken with wetting agents as russetting can occur. Both spray and dip methods can be applied. The correct timing of the application is critical for the success of the treatment.
Sultanas (seedless) Berry thinning (1)	1 tablet / 100 ℓ water	First application: When 50 % capfall has occurred in at least half of all bunches (half of all the bunches are 50 % in bloom).
Berry thinning (2)	1 tablet / 100 ℓ water	Second application: When 80 % capfall has occurred in at least half of all the bunches (half of all the bunches are 80 % in bloom).
Lower Orange River area To enlarge berries	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	First application: Apply when berries have set and 50 % are ± 4 to 5 mm in diameter.
	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	Second application: Apply when 75 % of berries are ± 4 to 5 mm in diameter.
	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	Third application: Apply when 100 % of berries are ± 4 to 5 mm in diameter.
All other areas To enlarge berries	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	First application: Apply when berries have set and are 5 mm in diameter.
	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	Second application: Apply 7 days after first application. (Application must not occur later than the pea-berry stage).

Crop / Use	Dosage Rate	Remarks
Grapes (continue) Flame Seedless Lower Orange River area Berry thinning	1 tablet / 200 ℓ water (0.5 tablet / 100 ℓ water)	Flame Seedless has naturally straggly bunches and therefore thinning may not be required in some instances. Apply during flowering when 80 % in full bloom.
Flame Seedless Lower Orange River area To enlarge berries (1)	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	First Application: Apply when berries are ± 7 to 8 mm in diameter.
To enlarge berries (2)	3 tablets / 100 ℓ water PLUS non-ionic wetting agent	Second Application: Apply when berries are ± 8 to 9 mm in diameter.
Waltham Cross Increase berry size and obtain a more even size berry	1 tablet / 200 ℓ water (0.5 tablet / 100 ℓ water) PLUS non-ionic wetting agent	Apply when Waltham Cross berries is 4 to 5 mm in diameter. Fill a container, which is large enough to hold one bunch with FALGRO® TABLETS solution and dip the entire bunch. Ensure that all berries are well wetted.
Hops To improve yield by enhancing growth in spring and for a shorter and more even hop training period	1 tablet / 100 ℓ water PLUS non-ionic wetting agent	Apply as a full cover spray during early spring; during September to middle October.
Mangoes To inhibit flowering of young trees	10 tablets / 100 ℓ water PLUS non-ionic wetting agent	Before FALGRO® TABLETS are applied, break off all flower panicles. Apply as a full cover spray during the last week of July or during the first week of August.
Potatoes Uncut tubers To improve sprouting of potato tubers	Winter planting areas only. Variable results occur in the summer rainfall regions. 1 tablet / 200 ℓ water (0.5 tablet / 100 ℓ water)	Dip tubers for a period of 10 minutes in FALGRO® TABLETS solution. Leave in a cool dry place and allow tubers to dry. Plant as soon as first eye movement is observed. Change to a fresh solution after dipping sufficient tubers for planting one hectare. Do not dip cut tubers in this solution, as the result will be unsatisfactory.
Cut tubers To improve sprouting of potato tubers	1 tablet / 1000 ℓ water	Dip tubers for a period of 10 minutes in FALGRO® TABLETS solution after cutting. Leave in a cool dry place and allow tubers to dry. Plant as soon as first eye movement is observed. Change to a fresh solution after dipping sufficient tubers for planting one hectare. Do not dip uncut tubers in this solution, as results will be unsatisfactory.
Forelle Pears To improve fruit set	1 tablet / 100 ℓ water	Apply as a full cover spray at 30 to 40 % flowering. Follow up with normal hand thinning 14 days after application.

Crop / Use	Dosage Rate	Remarks
Citrus		<ul style="list-style-type: none"> Apply as a full cover spray. Do not apply to orchards where Black spot has not been well controlled or where there is a high incidence of wind scar and Thrip markings. Do not apply to trees suffering from Root disease or generally neglected trees. Do not allow trees to go into severe wilt. If a summer oil has been sprayed on citrus, a minimum of 14 days must elapse before FALGRO® TABLETS may be applied. Oil may be applied 5 to 10 days after a FALGRO® TABLETS application. Do not apply 2,4-D Ester to citrus destined for export before consulting the export guidelines.
Clementines To increase fruit set	0.5 to 2 tablets / 200 ℓ water PLUS non-ionic wetting agent	Apply as a full cover spray from 65 to 75 % petal drop to 4 weeks after 100 % petal drop. With later applications applied, less influence on the fruit set reaction can be expected. The history of an orchard will influence FALGRO® TABLETS applications. Decide on the yield/fruit size balance before application. An excessive quantity of fruit will result in small fruit and induce alternate bearing. Get technical advice before applications are performed.
Grapefruit Extension of on-tree storage period (delay of harvesting), larger and firmer fruit	1 tablet / 100 ℓ water PLUS 2 ml 2,4-D Ester PLUS non-ionic wetting agent	Apply as a full cover spray at colour break-transparency (colour T5) or up to 3 weeks after colour break. Do not apply before colour break. Apply only to healthy trees where Black spot and Thrips are under control and no excessive wind scar occurs. The average fruit size must not be too large. The average diameter should exceed 89 mm at the time of spraying. Do not over-irrigate grapefruit trees intended for hanging.
Lemons To delay maturity (harvesting), thus to protect the crop during the decay prone period, fewer pickings and for larger fruits	1 tablet / 100 ℓ water PLUS 2 ml 2,4-D Ester PLUS non-ionic wetting agent	Apply as a full cover spray 5 to 8 weeks before normal picking starts. Apply only to trees free of Black spot or other blemishes.
Navels To reduce rind creasing in Navel oranges on Trifoliolate or Trifoliolate hybrid rootstock	2 tablets / 100 ℓ water PLUS non-ionic wetting agent	Apply as a full cover spray in January to February but not later than the last week of February. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cull on the total crop.
To reduce rind creasing in Navel Oranges on Rough lemon rootstock	1 tablet / 100 ℓ water PLUS non-ionic wetting agent	Apply as a full cover spray in January to February but not later than the last week of February. Apply only to healthy trees where Thrip blemish and windscars do not exceed 20 % cull on the total crop.
	OR 1 tablet / 100 ℓ water PLUS 2 ml 2,4-D Ester 500 PLUS non-ionic wetting agent	OR Apply as a full cover spray in May after full yellow colour has developed (minimum colour T3). Follow this treatment up with a 1 % urea (low biuret) at budbreak to avoid reduced flowering in the new season. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cull on the total crop.

Crop / Use	Dosage Rate	Remarks
Citrus (continue) Valencia Oranges To reduce rind creasing in Valencia oranges on all rootstocks	2 tablets / 100 ℓ water PLUS non-ionic wetting agent	Apply as a full cover spray in January to February but not later than the last week of February. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cull on the total crop.
	OR 1 tablet / 100 ℓ water PLUS non-ionic wetting agent	OR Apply as a full cover spray after the fruit has developed a full yellow colour (minimum colour T3). Do not apply before middle June. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cull on the total crop.
	OR 1 tablet / 200 ℓ water (0.5 tablet / 100 ℓ water) PLUS non-ionic wetting agent	OR Apply as a full cover spray when the fruit has developed a light greenish colour (T5 to T4, minimum colour T5). Do not apply before middle June. Apply only to healthy trees where Thrip blemish and wind scars do not exceed 20 % cull on the total crop.

FALGRO® is the registered trademark of / is die geregistreerde handelsmerk van **Fine Agrochemicals Limited.**

PRODUCT WARRANTY CONDITIONS AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications indicated on this label. FINE disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance or other unintended consequences may result because of factors such as abnormal weather conditions, presence of other materials, the manner of application, use of the Product other than in strict accordance with this label's instructions or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, the Buyer shall assume all such risks.

FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product. FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product.

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ, United Kingdom.