

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Fungicide



FEMIDA 500 SC

Reg. No. L 9638 Act/Wet No. 36 of/van 1947

2: 19/12/2019-Juni 2020

A suspension concentrate fungicide with systemic action for the control of fungal diseases on crops as listed.

'n Suspensie konsentraat swamdoder met sistemiese werking vir die beheer van swamsiektes op gewasse soos aangedui.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

fenhexamid (hydroxylanilide) **500 g/l** fenheksamid (hidroksie-anilied)

FRAC FUNGICIDE GROUP CODE **17** FRAC SWAMDODER GROEPKODE



villa

Registration holder / Registrasiehouer:

Universal Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07

PO Box / Posbus 801, Kempton Park, 1620

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: 3082

Willow Set & Print 011 394-4486



**CAUTION
VERSIGTIG**



GEBRUIKSAANWYSYNGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

FEMIDA 500 SC

Reg. No. L 9638 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: 17

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

fenhexamid (hydroxyanilide) / fenheksamied (hidroksie-anilied)..... 500 g/l

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION. (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620, Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS**

Allow the following number of days between the last application and harvest or grazing of the crops as listed below:	
Strawberries	1 day
Table grapes	3 days
Wine grapes	7 days

NOTE

ABOVE-MENTIONED WITHHOLDING PERIODS REFER TO COMPLIANCE WITH LOCAL MAXIMUM RESIDUE LIMITS (MRL'S). HOWEVER, IT IS IMPORTANT TO NOTE THAT IMPORT TOLERANCES OF OTHER COUNTRIES MIGHT POSSIBLY BE EXCEEDED. IF THE TREATED CROP WILL BE EXPORTED, FIRST CONSULT THE RELEVANT IMPORTER OR EXPORTING BODY REGARDING THE USE OF THIS PRODUCT, MRL'S AND RECOMMENDED WITHHOLDING PERIODS.

- Handle with care.
- May be harmful if swallowed.
- May cause damage if in contact with eyes.
- Toxic to aquatic organisms with long lasting effects.
- Store the product in the original container in a cool, dry, well-ventilated place away from food, seed, feedstuff and other agricultural chemicals.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other products not indicated on the label and the occurrence of resistance of the disease to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

- Avoid inhalation of the spray mist.
- Avoid eye and skin contact.
- Wear a face shield, rubber gloves and boots when handling the concentrate, preparing the spray mixture and during application.
- Wash with soap and water after use and accidental skin contact.
- In case of eye contamination, rinse the eyes thoroughly with plenty of clean water for at least 15 minutes, holding the eyelid(s) open. Obtain medical assistance immediately.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke whilst mixing and applying the product or before washing hands and face and change of clothing.
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment or to nearby water sources.
- Thoroughly clean spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.

- **TRIPLE RINSE** empty containers in the following manner. Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening and dispose of it in a safe way.
- **Do not** re-use the empty container for any other purpose.
- Prevent contamination of food, feed, drinking water and eating utensils.

RESISTANCE WARNING

FEMIDA 500 SC is a group code 17 fungicide. Any fungus population may contain individuals naturally resistant to **FEMIDA 500 SC** and other group code 17 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **FEMIDA 500 SC** or any other group code 17 fungicide.

To delay fungicide resistance:

- avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes,
- integrate other control methods (chemical, cultural, biological) into disease control programmes.

For specific information on resistance management contact the registration holder of this product.

USE RESTRICTIONS

- Allow seven (7) days between the application of **FEMIDA 500 SC** and a dust application or products not listed above and do not apply **FEMIDA 500 SC** in combination with wetter/stickers or as a tank mixture with trace or macro element sprays in grapes.

DIRECTIONS FOR USE: Only use as indicated.

Compatibility:

- Refer to “**USE RESTRICTIONS**” above.
- Although **FEMIDA 500 SC** seems to be compatible with most other agrochemical products, the compatibility of **FEMIDA 500 SC** with other products cannot be guaranteed. A physical compatibility test must always be performed before such a tank mixture is sprayed. Water quality may influence the compatibility of **FEMIDA 500 SC**.
- When **FEMIDA 500 SC** is used in conjunction with any other agricultural remedy, adhere to all **WARNINGS, PRECAUTIONS** and **DIRECTIONS FOR USE** mentioned on that label.

Mixing instructions:

- Shake the **FEMIDA 500 SC** container well before use.
- Half fill the spray tank with clean water.
- Premix the required quantity of **FEMIDA 500 SC** with at least 10 litres of water.
- Add the premix to the water in the spray tank.
- Agitate well to obtain a homogeneous emulsion.
- Fill the spray tank with water to the required level while maintaining agitation, to ensure thorough mixing.
- Maintain agitation while spraying.
- If another product is to be mixed with **FEMIDA 500 SC**, the required quantity of the other product should be premixed, prior to slowly adding to the spray mixture.
- Do not leave the prepared spray mixture in the spray tank for any length of time, e.g. overnight.

Application instructions:

FEMIDA 500 SC can be applied with conventional high-volume spray equipment. It is important that the sprayer will be in good working order and correctly calibrated prior to application, to ensure that the correct delivery rate and even distribution of the spray mixture is obtained.

APPLICATION RATES**Remarks regarding resistance management in berries:**

- **Do not** exceed three (3) **FEMIDA 500 SC** applications per season per crop. If three (3) **FEMIDA 500 SC** treatments are applied, it must **not** be applied consecutively.
- Precede/alternate and follow the treatments in the spray programme with registered fungicides with a different mode of action.

Remarks regarding resistance management in vines:

- **Do not** exceed three (3) **FEMIDA 500 SC** applications per season per vineyard. If three (3) **FEMIDA 500 SC** treatments are applied, it must **not** be applied consecutively.
- Precede/alternate and follow the treatments in the spray programme with registered fungicides with a different mode of action e.g. **Source 400 SC**, **Iprodione 500 SC**, **Chant 500 WDG**, **Volume 500 WDG**.

Crop and disease	Dosage rate	Remarks
Strawberries <i>Botrytis cinerea</i>	75 ml / 100 l water OR 7.5 ml / 10 l water	Apply during flowering as a full cover spray at seven (7) day intervals. A preventative spray programme is recommended for the control of <i>Botrytis</i> which should commence at the onset of flowering. Do not apply more than three (3) FEMIDA 500 SC treatments per season. Alternate applications of FEMIDA 500 SC with registered <i>Botrytis</i> controlling fungicides with different modes of action.
Table Grapes <i>Botrytis cinerea</i>	75 ml / 100 l water (750 ml to 1125 ml/ha)	Apply at flowering and the pea size stage of the berries as a high or low volume spray (refer NOTES below). FEMIDA 500 SC can also be applied at the start of ripening and shortly before harvest as a low volume spray application in 500 to 700 litres water per hectare with a suitably calibrated mistblower, delivering a droplet spectrum with a VMD of 150 to 200 µm. Do not spray past the point of run-off on the berries. Do not apply more than three (3) FEMIDA 500 SC treatments per season. For the control of <i>Botrytis</i> at growth stages not mentioned, use registered <i>Botrytis</i> controlling fungicides with different modes of action.
Wine Grapes <i>Botrytis cinerea</i>	75 ml / 100 l water (750 ml to 1125 ml/ha)	Treatments can be applied at full blossom stage, at pea size stage of the berries, at the start of ripening (<i>véraison</i>) or shortly before harvest. Do not apply more than three (3) FEMIDA 500 SC treatments per season. For the control of <i>Botrytis</i> at growth stages not mentioned, use alternative registered <i>Botrytis</i> controlling fungicides with different modes of action.

NOTE**1. High volume application:**

Spray to the point of run-off ensuring even coverage. Depending on the leaf area present, 750 to 1500 litres spray mixture should be applied per hectare as follows:

Growth stage	Water requirements per hectare as recommended according to the different growth stages of a tree:	
	Table grapes	Wine grapes
Blossom to pea-size stage of the berries	1000 to 1500	750 to 1000
Berries at pea-size stage up to harvest	Refer to NOTE 2 Low volume application	1000 to 1500

2. Low volume application:

- **FEMIDA 500 SC** can be applied as a low volume application. The volume of product required per hectare should be calculated from the corresponding high-volume application rate.
- The concentration of the spray mix should not exceed four times (4X) the highest concentration of a high-volume application (i.e. highest dosage in the lowest volume of water of the high volume range indicated above).
- Wine grapes: Apply in at least 250 litres of water per hectare and ensure even coverage.
- Table grapes: Apply in 500 to 700 litres water per hectare with a suitably calibrated mistblower at the start of ripening or shortly before harvest. Apply a droplet spectrum with a VMD of 150 to 200 µm up to the point of run-off and ensure even coverage.
- Do not spray past the point of run-off on the grape berries. Calculate the volume of product required per hectare from the high-volume application rate applicable to the particular vineyard at that growth stage.

The following products mentioned in this label can be replaced with the following equivalent products:

- **SOURCE 400 SC** (L 9283) = **SUPPORT 400 SC** (L 9282) (**Pyrimethanil**),
- **IPRODIONE 500 SC** (L 8459) = **TAURUS 500 SC** (L 8499 / N-AR 1137) (**Iprodione**),
- **CHANT 500 WDG** (L 8958) = **CHOIR 500 WDG** (L 8957) (**Cyprodinil**) and
- **VOLUME 500 WDG** (L 9261) = **CLOUD 500 WDG** (L 9260) (**Boscalid**).

SOURCE 400 SC, IPRODIONE 500 SC, CHANT 500 WDG and/en **VOLUME 500 WDG** are registered products of / is geregistreerde produkte van **UNIVERSAL CROP PROTECTION (PTY) LTD.**

SUPPORT 400 SC, TAURUS 500 SC, CHOIR 500 WDG and/en **CLOUD 500 WDG** are registered products of / is geregistreerde produkte van **VILLA CROP PROTECTION (PTY) LTD.**