

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Fungicide



FIGURE 500 SC

Reg. No. L 9920 Act/Wet No. 36 of/van 1947

1: 12/10/2015 – June 2017

A suspension concentrate organic based fungicide for the preventive control of plant diseases in crops as listed.

in Suspensie konsentraat organies-gebaseerde swamdoder vir die voorkomende beheer van plantsiektes in gewasse soos aangedui.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

folpet (phtalimide) **500 g/l** folpet (ptalimied)

FRAC FUNGICIDE GROUP CODE **M4** FRAC SWAMDODER GROEPKODE



villa

Registration holder / Registrasiehouer:
Universal Crop Protection (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07
PO Box / Posbus 801, Kempton Park, 1620
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: 3082

Willow Set & Print 011 394-4486



CAUTION
VERSIGTIG



GEBRUIKSAANWYINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

DIRECTIONS FOR USE ENCLOSED

Batch number:
Lotnommer:

FIGURE 500 SC

Reg. No. L 9920 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: M4

ACTIVE INGREDIENT/ AKTIEWE BESTANDDEEL:

folpet (phtalimide)

.....500 g/l

folpet (ptalimied)

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

PO Box / Posbus 801,

KEMPTON PARK, 1620, Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS****Withholding periods:**

Minimum time between the last application and harvest and/or grazing:	
Table grapes	Not after the pre-blossom stage.
Wine grapes	14 days

NOTE

ABOVE-MENTIONED WITHHOLDING PERIODS REFER TO COMPLIANCE WITH LOCAL MAXIMUM RESIDUE LIMITS (MRL'S). HOWEVER, IT IS IMPORTANT TO NOTE THAT IMPORT TOLERANCES OF OTHER COUNTRIES MIGHT POSSIBLY BE EXCEEDED. IF THE TREATED CROP WILL BE EXPORTED, FIRST CONSULT THE RELEVANT IMPORTER OR EXPORTING BODY REGARDING THE USE OF THIS PRODUCT, MRL'S AND RECOMMENDED WITHHOLDING PERIODS.

- Handle with care.
- Eye irritant and possible skin sensitizer upon contact.
- Toxic to fish.
- Store in a cool, dry place away from food, feeds, fertilizers and other agricultural remedies.
- Keep out of reach of children, animals and uninformed persons.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Avoid eye and skin contact and inhalation of spray mist.
- Wear rubber gloves and a face shield when handling the concentrate and application of the spray mixture.
- In case of eye contact, flush out at once.
- In case of skin contact, wash off at once.
- In case of spillage of product on clothes, change and wash immediately.
- Do not eat, drink or smoke whilst mixing, applying the product or before washing hands and face or change of clothing.
- Avoid drift of spray onto other crops, grazing, rivers, dams and areas not under treatment.
- Thoroughly clean the spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- Invert the empty container over the spray or mixing tank to make sure the container is empty before destroying the container.

- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

RESISTANCE WARNING

FIGURE 500 SC is a group code M4 fungicide. Any fungus population may contain individuals naturally resistant to **FIGURE 500 SC** and other group code M4 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly and exclusively in programs. These resistant fungi may not be controlled by **FIGURE 500 SC** or any other group code M4 fungicides.

To delay fungicide resistance:

- avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate with products from different fungicide group codes,
- integrate other control methods (chemical, cultural, biological) into insect control programmes.

Since the presence of resistant fungi is difficult to detect prior to fungicide application, it is of the utmost importance that treated areas be inspected at regular intervals, to identify the occurrence of the fungicide resistant fungi timeously. The registration holder will not accept liability for performance failures in the event of a build-up of resistant fungi resulting from inadequate resistance management practices as indicated on the label.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed

Compatibility:

- **FIGURE 500 SC** can be applied in tank mixtures with the most commonly used insecticides and fungicides with the exception of **Lime, Lime Sulphur** and **Bordeaux** mixture. It should also **NOT** be applied with any other *Botrytis* remedy in wine and table grapes.
- When **FIGURE 500 SC** is used in conjunction with any other agricultural remedy, adhere to all **WARNINGS, PRECAUTIONS** and **DIRECTIONS FOR USE** mentioned on that label.
- A physical compatibility test must always be performed before such a tank mixture is sprayed. It is also recommended to do a small scale test application first before a large scale commercial treatment of such mixtures to confirm crop safety.
- **Do not apply more than 4 applications per season.**

Mixing instructions:

- **Shake the container thoroughly before tank mixture is prepared.**
- Half fill the spray tank with clean water and engage agitation.
- When the spray or pre-mix tank is half full with water, add **FIGURE 500 SC** slowly and agitate until completely dispersed.
- Agitation must be continued throughout the mixing and spraying operation. Re-suspend the spray mixture if agitation was suspended for longer than 5 minutes.
- If any other product is to be mixed with **FIGURE 500 SC**, pre-mix the required volume of product in a similar fashion. Add the pre-mixed product(s) in the following sequence (as applicable): Powder/granular formulation (e.g. WP/WDG), Suspension Concentrate (SC), Emulsifiable Concentrate/emulsion (EC/EW).
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

Application:

- Apply as a preventative application and ensure thorough coverage of both upper and lower leaf surfaces.
- **FIGURE 500 SC** can be applied with conventional high volume spray equipment. It is important that the sprayer be in good working order and correctly calibrated prior to application, to ensure that the correct delivery rate and even distribution of the spray mixture is obtained.

APPLICATION RATES

Crop / Pest	Dosage rate	Remarks
FIGURE 500 SC IS ALWAYS RECOMMENDED IN AN INTEGRATED DISEASE MANAGEMENT PROGRAMME. IN ORDER TO ACHIEVE CONSISTENT AND RELIABLE DISEASE CONTROL, PREVENTIVE FUNGICIDE APPLICATIONS ARE CRITICALLY IMPORTANT.		
<u>Wine grapes</u> Deadarm (<i>Phomopsis viticola</i>) Botrytis rot (<i>Botrytis cinerea</i>) Downy mildew (<i>Plasmopora viticola</i>)	200 ml / 100 ℓ water	Commence with treatment at bud swell. Repeat with 7 day intervals if climatological conditions promote disease development. Do not apply more than four (4) applications per season. <u>Pre-blossom:</u> Apply first treatment when shoots are 10 cm long and repeat at 7 to 10 day intervals depending on weather conditions. <u>Blossom to 80 % petal drop:</u> Only apply when the blossoming period lasts longer than 2 weeks. <u>Post blossom:</u> Apply twice or more at 7 to 14 day intervals depending on weather conditions. Do not apply more than four (4) applications per season.
<u>Table grapes</u> Deadarm (<i>Phomopsis viticola</i>) Downy mildew (<i>Plasmopora viticola</i>)	200 ml / 100 ℓ water	Recommended only until pre-blossom stage. Apply first treatment at bud swell and repeat at 7 day intervals. Do not apply more than four (4) applications per season.