

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Fungicide



FISSION 520 SC

Reg. No. L 9756 Act/Wet No. 36 of/van 1947

2: 10/9/2018 – Sept2018

A suspension concentrate systemic fungicide for the control of certain fungal diseases on crops as listed.

'n Suspensie konsentraat sistemiese swamdoder vir die beheer van sekere swamsiektes op gewasse soos gelys.

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE

pyrimethanil (aniline derivative) **400 g/ℓ** pirimetanil (anilien derivaat)
trifloxystrobin (strobilurine) **120 g/ℓ** trifloksistrobien (strobilurien)

FRAC FUNGICIDE GROUP CODE **9** FRAC SWAMDODER GROEPKODE

FRAC FUNGICIDE GROUP CODE **11** FRAC SWAMDODER GROEPKODE

kg

villa

Registration holder / Registrasiehouer:

Universal Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07

PO Box / Posbus 801, Kempton Park, 1620

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: 3082

Willow Set & Print 011 394-4486



CAUTION
VERSIGTIG



GEBRUIKSAANWYSLINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

DIRECTIONS FOR USE ENCLOSED

Batch number:
Lotnommer:

FISSION 520 SC

Reg. No. L 9756 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: M9 & 11

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

pyrimethanil (aniline derivative) / pyrimethanil (anilien derivaat)400 g/l

trifloxystrobin (strobilurine) / trifloksistrobien (strobilurien)120 g/l

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620

Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS****Withholding periods:**

| Allow the following number of days between the last application and harvest of the following crops: | |
|--|---------|
| Apples | 35 days |
| Pears | 35 days |
| Potatoes | 14 days |

NOTES

THE MAXIMUM RESIDUE (MRL'S) LEVELS WILL NOT BE EXCEEDED, PROVIDED APPLICATIONS ARE APPLIED AS HEREWITH PRESCRIBED. WHEN PRODUCTION IS DESTINED FOR THE EXPORT MARKET, THE EXPORT AGENCY OF CONCERN MUST BE CONTACTED, PRIOR TO THE USE OF FISSION 520 SC IN A SPRAY PROGRAMME.

TO PREVENT THAT PERMISSABLE MRL'S ARE EXCEEDED, DO NOT APPLY OTHER TRIFLOXYSTROBIN OR PYRIMETHANIL CONTAINING PRODUCTS IN THE SAME SEASON'S SPRAY PROGRAMME.

- Handle with care.
- May cause skin irritation.
- Toxic to fish and other aquatic organisms. Do not contaminate dams, rivers, lakes or any other water sources.
- Store the product in the original container and always keep the container tightly closed.
- Store in a cool (preferably below 30°C), dry, well-ventilated place, away from food, seed and animal feeds.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the diseases against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Avoid inhalation of the spray mist.
- Avoid prolonged contact of product with skin.
- Wear rubber gloves, a face shield and boots when handling the concentrate.
- Wash contaminated clothes daily.
- Wash thoroughly with soap and water after application and accidental skin contact.
- In case of eye contamination, rinse the eyes thoroughly with plenty of clean, cold water for at least 20 minutes, holding the eyelid(s) open. If irritation persists obtain medical aid immediately.

- Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and change of clothing.
- Prevent spray drift onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean the spraying equipment directly after use by rinsing with clean water and dispose of wash water where it will not contaminate crops, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feed, drinking water and eating utensils.

RESISTANCE WARNING

FISSION 520 SC contains active ingredients of FRAC group codes 9 & 11. Any fungus population may contain individuals naturally resistant to **FISSION 520 SC** and other group code 9 & 11 fungicides. The resistant individuals may eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **FISSION 520 SC** or any other group code 9 & 11 fungicides.

To delay fungicide resistance:

- avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes,
- for tank mixing or alternation with products in fungicide group codes 9 and 11, refer to applicable individual product labels,
- integrate other control methods (chemical, cultural, biological) into disease control programmes.

For specific information on resistance management, contact the registration holder of this product.

Additional notes on resistance management:

For the control of Apple & Pear scab as well as Powdery mildew on apples, it is essential to adhere to the following FRAC (Fungicide Resistance Action Committee) guidelines.

- Apply QoI (strobilurines) fungicides according to manufacturer's recommendations for the target disease (or complex) at the specific crop growth stage indicated and adapted to size of trees. Effective disease management is a critical parameter in delaying the build-up of resistant pathogen populations.
- QoI fungicides must only be applied in mixtures with partners contributing to the effective control of the target pathogens.
- Apply QoI fungicides preventively. Under high disease pressure the spray interval should not exceed 7 to 10 days.
- Apply a maximum of 3 QoI containing sprays per crop per season. A maximum of 4 QoI fungicide applications may be used where twelve (12) or more applications are made per crop per season.
- A maximum of two (2) consecutive QoI fungicide sprays is preferred. Where field performance was adversely affected apply QoI containing fungicides in mixtures in strict alternation with fungicides from a different group.

Refer to the FRAC Website for latest guidelines: (<http://www.frac.info/frac/index.htm>).

USE RESTRICTIONS

- **Do not** apply more than 3 treatments of **FISSION 520 SC** per season.

DIRECTIONS FOR USE: Only use as directed.

General information:

- Apply **FISSION 520 SC** primarily as a preventative application.
- Use alternative registered fungicides with a mode of action that differs from strobilurines / Group Code 11 (QoI) related fungicides.
- The last fungicide treatment of the season must not be a strobilurine related fungicide
- Use **FISSION 520 SC** as part of an IPM and Good Agricultural Practice (GAP) programme.

Compatibility:

- **Do not** apply in mixtures with aluminium containing products.

- The compatibility of **FISSION 520 SC** in a tank mixture with other products is influenced by the formulations of these products, as well as quality of the spray water. It is recommended that a physical compatibility test always be performed prior to application as the formulation of other products can be altered and the quality of water can vary from area to area.

Mixing instructions:

- **Shake container well before opening.**
- Half-fill the mixing tank with clean water.
- Premix the required quantity of **FISSION 520 SC** in at least 10 litres of water in another container.
- Add the **FISSION 520 SC** mixture into the mixing tank and fill the mixing tank up to the required volume of water.
- If another product is to be added to the mixing tank, premix the required amount of product with at least 10 litres of water, before adding to the mixing tank.
- Ensure continuous agitation during mixing and application.
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

Application instructions:

- Apply **FISSION 520 SC** with correctly calibrated spray equipment that is in good working order.
- The spray volume required for a specific growth stage must be determined accurately according to the Tree-Row-Volume method as described by Unrath (Deciduous Fruit Grower, November 1986). The application volume rate per hectare is then determined according to the spray volume calculated for high volume application per hectare. Refer to the table below.

APPLICATION RATES

| <i>Crop / Disease</i> | <i>Dosage rate</i> | <i>Remarks</i> |
|--|--------------------|---|
| Potatoes Early blight <i>(Alternaria solani)</i> Brown spot <i>(Alternaria alternata)</i> | 770 ml / ha | <p>Refer to “Resistance Management, Directions for use and Crop recommendations” above.</p> <p>Apply preventively as part of a 7-day spray programme.</p> <p>Start with a weekly program application one (1) week after the potatoes have emerged. Use alternative effective fungicides as registered on potatoes up to week 5 in a 7 day spray interval. Apply FISSION 520 SC at 14-day intervals during week 5, 7 and 9 alternated with either Mancozeb containing fungicides or Mission 720 SC during week 6, 8 and 10. Revert back to alternative effective fungicides in a 7-day spray program until such time when disease control is not required anymore.</p> <p>Do not apply more than three (3) FISSION 520 SC applications per growing season.</p> <p><u>Ground application:</u> At least 500 litres water per hectare. <u>Aerial application:</u> 30 to 40 litres water per hectare.</p> |

| Crop / Disease | Dosage rate | Remarks |
|---|---|---|
| Apples Scab (<i>Venturia inaequalis</i>) Powdery mildew (<i>Podosphaera leucotricha</i>) | 40 ml / 100 l water | Refer to “Resistance Management and Application instructions” above. Apply preventively as part of a spray programme. |
| Pears Scab (<i>Venturia pirina</i>) | | Start with the applications at the green-tip stage and apply at weekly intervals until 75 % petal fall. After 75 % petal fall, the spray programme intervals can be extended to 14 days. However, should weather conditions favour disease development during this time, revert back to weekly intervals. FISSION 520 SC can be included during any time of the spray programme. However, it is recommended to apply FISSION 520 SC between flowering and early fruit set for optimum disease control. |
| IMPORTANT Use the Tree Row Volume (TRV) formula below to ensure that the correct amount of fungicide is applied per hectare. This is calculated as follows: <u>High volume rate:</u> The water to be applied per hectare must be calculated according to the TRV formula: $l / ha = \frac{\text{Tree height} \times \text{Tree diameter} \times 937}{\text{Row width}}$ The TRV calculation is the maximum water volume required when the trees are in full leaf. Use the water volume required per hectare to calculate the amount of FISSION 520 SC required per hectare according to the various growth stages of a tree during the season. | | |
| Water requirements per hectare as recommended according to the different growth stages of a tree: | | |
| <u>Growth stage</u> | <u>% of the high volume water requirement / ha</u> | |
| From greentip to ± 30 % blossom | 60 % | |
| From ± full blossom to ± middle November | 80 % | |
| From beginning December | 100 % | |

Consult all product labels mentioned in this label for **WARNINGS, PRECAUTIONS, USE RESTRICTIONS** and **DIRECTIONS FOR USE**.

The following product mentioned in this label may be replaced with an equivalent product:

- **MISSION 720 SC** (L 8432 / N-AR 1298) = **ORONIL 720 SC** (L 8431).

ORONIL 720 SC is a registered product of / is 'n geregistreerde produk van
UNIVERSAL CROP PROTECTION (PTY) LTD.

MISSION 720 SC is a registered product of / is 'n geregistreerde produkte van
VILLA CROP PROTECTION (PTY) LTD.