

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

SHAKE THE BOTTLE WELL BEFORE USE.

Fungicide



FLUSILAZIM EXTRA 375 SC

Reg. No. L 9500 Act/Wet No. 36 of/van 1947

1: 11/3/2014 – July 2017

A suspension concentrate fungicide for the control of diseases on crops as indicated.

'n Suspensie konsentraat swamdoder vir die beheer van siektes op gewasse soos aangedui.

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE

flusilazole (triazole) **125 g/l** flusilasool (triasool)
carbendazim (benzimidazole) **250 g/l** karbendasim (bensimidasool)

FRAC FUNGICIDE GROUP CODE **1** FRAC SWAMDODER GROEPKODE
FRAC FUNGICIDE GROUP CODE **3** FRAC SWAMDODER GROEPKODE

5 l

villa

Registration holder / Registrasiehouer:
Universal Crop Protection (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 1983/008184/07
PO Box / Posbus 801, Kempton Park, 1620
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: 3082

Willow Set & Print 011 394-4486



HARMFUL
SKADELIK



GEBRUIKSAANWYSINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

DIRECTIONS FOR USE ENCLOSED

Batch number:
Lohnommer:

FLUSILAZIM EXTRA 375 SC

Reg. No. L 9500 Act/Wet No. 36 of/van 1947

FRAC FUNGICIDE GROUP CODE / FRAC SWAMDODER GROEPKODE: 3 & 1

ACTIVE INGREDIENTS / AKTIEWE BESTANDEDELE:

flusilazole (triazole) / flusilasool (triasool) 125 g/l
 carbendazim (benzimidazole) / karbendasim (bensimidasool)..... 250 g/l

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION. (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620, Tel. (011) 396 2233

HARMFUL**SKADELIK****WARNINGS**

Allow the following number of days between the last application and harvest or grazing of the crops listed below:	
Barley & Wheat	56 days
Groundnuts (fodder)	42 days
Soybeans	30 days
Soybeans (fodder)	42 days

NOTE

ABOVE-MENTIONED WITHHOLDING PERIODS REFER TO COMPLIANCE WITH LOCAL MAXIMUM RESIDUE LIMITS (MRL'S). HOWEVER, IT IS IMPORTANT TO NOTE THAT IMPORT TOLERANCES OF OTHER COUNTRIES MIGHT POSSIBLY BE EXCEEDED. IF THE TREATED CROP WILL BE EXPORTED, FIRST CONSULT THE RELEVANT IMPORTER OR EXPORTING BODY REGARDING THE USE OF THIS PRODUCT, MRL'S AND RECOMMENDED WITHHOLDING PERIODS.

- Do not graze treated barley and wheat fields.
- Handle with care.
- Harmful when swallowed, inhaled or absorbed through the skin.
- May cause skin and eye irritation.
- Toxic to fish and aquatic organisms.
- Do not contaminate dams, rivers, lakes or any other water sources.
- Store in a cool, dry, well-ventilated place in the original container, away from food and feedstuffs.
- Avoid storage temperatures below -5 °C and above 25 °C.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area within 1 day after treatment unless wearing protective clothing.
- **In case of poisoning call a doctor and make this label available to him/her.**

Aerial application:

Notify all inhabitants in the immediate vicinity of the area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate adjacent areas or water.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Avoid inhalation of the spray mist.
- Avoid contact with skin and eyes.
- Wear a face shield and rubber gloves when preparing the mixture and during application.

- Wash contaminated clothing after use.
- Wash with soap and water after use and after accidental skin contact.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face and change of clothing.
- Do not spray or allow drift of spray on to other crops, grazing, rivers, dams and areas not under treatment or to nearby water sources.
- Clean the spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of a third of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the container by perforation and flattening and dispose of it in a safe way.
- **Do not** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

SYMPTOMS OF HUMAN POISONING

Prolonged or repeated exposure may cause eye damage or irritate the respiratory tract and eyes, and may cause headaches and dizziness. Some individuals may develop an allergic response.

FIRST AID TREATMENT

- **Inhalation:** Remove the patient from the source of contamination to fresh air. Obtain medical attention if irritation persists.
- **Skin contact:** Wash skin gently and thoroughly with cold water and non-abrasive soap. Obtain medical attention if irritation persists.
- **Eye contact:** Immediately irrigate eyes with clean water for at least 20 minutes. Obtain medical attention if irritation persists.
- **Ingestion:** Do not induce vomiting or give anything by mouth. Obtain medical attention immediately. If the person is alert, administer 1 to 2 glasses of water and, if possible, 30 to 50 g of activated charcoal.

NOTE TO PHYSICIAN

There is no specific antidote available. Treat symptomatically. Consider gastric lavage avoiding aspiration. Do not give ephedrine or related drugs.

RESISTANCE WARNING

FLUSILAZIM EXTRA 375 SC is a mixture of group codes 3 and 1 fungicides. Any fungus population may contain individuals naturally resistant to **FLUSILAZIM EXTRA 375 SC** and other group code 3 and 1 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **FLUSILAZIM EXTRA 375 SC** or any other group code 3 and 1 fungicides.

To delay fungicide resistance:

- avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes,
- for tank mixing or alternation with products in fungicide group code M, refer to applicable, individual product labels,
- integrate other control methods (chemical, cultural, biological) into disease control programmes.

For specific information on resistance management contact the registration holder of this product.

GENERAL PROPERTIES

- **FLUSILAZIM EXTRA 375 SC** is a mixture of systemic foliar fungicides containing **Flusilazole**, a DMI fungicide, and **Carbendazim**, an MBC fungicide, with both protective and post-infection curative action.
- The efficiency of **FLUSILAZIM EXTRA 375 SC** is not influenced by rain occurring 3 hours after application, due to rapid uptake of the active ingredients by plant tissue.

USE RESTRICTIONS

- It is not recommended to apply systemic products, such as **FLUSILAZIM EXTRA 375 SC**, when crops are under severe drought and/or fertility stress conditions. The uptake and activity of systemic compounds may be reduced under these conditions. Consult a representative or the distributor in the event of any uncertainty.

DIRECTIONS FOR USE: Use only as directed.**Compatibility:**

- **FLUSILAZIM EXTRA 375 SC** is compatible with **Link**, other Villa approved adjuvants and most commonly used fungicides, insecticides and foliar feeds normally used with the various crops.
- The compatibility of **FLUSILAZIM EXTRA 375 SC** with other products may be influenced by various factors. A physical compatibility test must always be performed prior to application of such a tank mixture.
- When **FLUSILAZIM EXTRA 375 SC** is used in combination with any other agricultural remedy, adhere to all **WARNINGS, PRECAUTIONS** and **DIRECTIONS FOR USE** mentioned on these labels.

Mixing instructions:

- Half fill the spray tank with clean water.
- Shake the **FLUSILAZIM EXTRA 375 SC** container well before use.
- Add the required amount of **FLUSILAZIM EXTRA 375 SC** to the water in the spray tank while stirring.
- If any other product is to be mixed with **FLUSILAZIM EXTRA 375 SC**, the required volume of this product must be pre-mixed in a mixing tank with 10 litres water. When a wettable powder is to be added, cream in advance.
- Agitate the water in the spray tank and then add the product(s) to the tank in the following sequence (as applicable): acidifier/buffer or adjuvant, suspension concentrate, water soluble concentrate, emulsifiable concentrate.
- Fill the spray tank with water to the required level while maintaining agitation, to ensure thorough mixing.
- Maintain agitation while spraying.
- Prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

APPLICATION**Ground application:**

FLUSILAZIM EXTRA 375 SC can be applied with conventional high volume spray equipment. Calibrate the apparatus before application, to ensure that the correct dosage is applied. The distribution of the spray solution must be even throughout the target area.

Aerial application:

Aerial application of **FLUSILAZIM EXTRA 375 SC** may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- Volume: A spray mixture volume of 30 litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.
- Droplet coverage: 25 to 35 droplets per cm² must be recovered at the target area.
- Droplet size: A droplet spectrum with a VMD of 280 to 300 micron is recommended. Limit the production of fine droplets less than 150 micron (high drift and evaporation potential) to a minimum.
- Flying height: Maintain the height of the spray boom at 3 to 4 meters above the target. Do not spray when aircraft dives, is in a climb or when banking.
- Use suitable atomising equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8°C.
- Stop spraying if the wind speed exceeds 15 km per hour.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage),
 - b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

APPLICATION RATES

Crop and Disease	Dosage rate	Remarks
Wheat & Barley Eye spot (<i>Pseudocercospora herpotrichoides</i>)	700 ml (Ground application) OR 840 ml (Aerial application)	Apply between the 5- and 7-leaf crop stages. Ground application in at least 300 litres water per hectare and Aerial application in at least 30 litres water per hectare.
Groundnuts Leafspot (<i>Cercospora arachidicola</i>) (<i>Cercosporidium personatum</i>) Leaf blotch (<i>Phoma arachidicola</i>) Botrytis (<i>Botrytis cinerea</i>)	500 ml (Ground application) OR 675 ml (Aerial application)	Ground application in at least 500 litres water per hectare and Aerial application in at least 30 litres water per hectare. Aerial application on dry land groundnuts only. Apply as a full cover spray, starting when the first symptoms of disease appear. Repeat at 10- to 14-day intervals up to 2 weeks before lifting of the groundnuts. Do not apply more than 6 applications per season. Allow at least 42 days to lapse between the last application and use of straw for feeding.
Rust (<i>Puccinia arachidis</i>)	1000 ml in 500 litres water per ha	Apply as discussed for Leafspot and Blotch disease. Ground application only.
Soybeans	Ground application in at least 500 litres water per hectare and Aerial application in at least 40 litres water per hectare. Ensure thorough coverage of the foliage.	
Ascochyta leafspot (<i>Ascochyta</i> spp.)	600 ml (Ground application)	Apply when the first signs of disease appear on the foliage and repeat 10 to 14 days later if climatic conditions favour disease development.
Rust (<i>Phakopsora pachyrhizi</i>)	OR 750 ml (Aerial application) PLUS Link (50 ml / 100 l water)	Apply at the onset of flowering or first signs of disease around early flowering. Ensure thorough coverage of the whole plant. If the first application is applied during flowering or at early pod-fill, a follow-up application 3 to 4 weeks later is recommended. If weather conditions favour disease development (high humidity and temperatures of 15 to 28 °C) use the shorter interval. A single application will be sufficient should the disease appear late in the growing season. Allow at least 42 days between the last application and use of hay for feeding. Do not exceed 2 applications per season.

NOTES

- Diseases are best controlled if fungicide applications are performed preventively. Applications should therefore be performed before infection takes place or when disease incidence is low.
- Fungicides should not be applied when plants are under severe drought stress or when high infection pressure is present.

The following products mentioned in this label may be replaced with an equivalent products:

- **LINK** (L 8675) = **DIRECT** (L 8680).

LINK and/en **DIRECT** are registered products of / is geregisteerde produkte van
VILLA CROP PROTECTION (PTY) LTD.