

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Micronutrient



MAX-IN[®] ZINC

Reg. No. B 5278 Act/Wet No. 36 of/van 1947

1: 09/02/2017 – Oct2019

A water-soluble foliar feed for use in a balanced fertilizer program for all crops requiring zinc.

'n Wateroplosbare blaarvoeding vir gebruik in 'n gebalanseerde misstof program op alle gewasse wat sink benodig.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

zinc / sink (Zn).....60 g/kg
SG: 1.2 at 20 °C

FERTILIZER GROUP/GROEP 2 MISSTOF



villa The Villa logo consists of the word "villa" in a blue, lowercase, sans-serif font, followed by a red diamond shape containing a white circle.

Registration holder / Registrasiehouer:
Winfield Solutions Registration Holdings (Pty) Ltd.
Co. Reg. No. / Mpy. Reg. Nr. 2015/312008/07
PO Box / Posbus 10413, Aston Manor, 1630
Tel: 011 396 2233
Website / Webblad: www.villacrop.co.za

UN Number: Not Regulated

GEBUIKSAANWYSINGS INGESLUIT

VERWYS NA BESONDERHEDE
GEDRUK OP HOUER/SAK

Date formulated:
Formuleringsdatum:

DIRECTIONS FOR USE ENCLOSED

REFER TO DETAILS PRINTED
ON CONTAINER/BAG

Batch number:
Lotnommer:

MAX-IN® ZINC

Reg. No. B 5278 Act No. 36 of 1947
 FERTILIZER GROUP/GROEP 2 MISSTOF

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL:

zinc / sink (Zn)60 g/kg

Registration holder / Registrasiehouer:

WINFIELD SOLUTIONS REGISTRATION HOLDINGS (PTY) LTD.

Co. Reg. No. 2015/312008/07

P.O. Box 10413, ASTON MANOR, 1630,

Tel. (011) 396 2233

DIRECTIONS FOR USE

- For use in a balanced fertilizer program for all crops requiring zinc.
- User should seek the advice of a certified crop advisor or trained agronomist.
- For specific crop recommendations in your area, consult your local agricultural professional.

MIXING INSTRUCTIONS

- Add water and agitate.
- Add **MAX-IN® ZINC**.
- At the end of the day, rinse out spray booms.

APPLICATION RATES

<i>Crop</i>	<i>Application rate</i>	<i>Remarks</i>
<u>Lucern</u>	2.25 to 4.5 t/ha	Apply MAX-IN® ZINC after each cutting. Applications should be made when sufficient re-growth has occurred.
<u>Cereal Crops</u>	2.25 to 4.5 t/ha	Apply when plants are between tillering and early joint stages. Later application may be needed; tissue samples should be used to determine later usages.
<u>Corn</u>	2.25 to 4.5 t/ha	Apply when corn is in the (V3 to V8) stage. Later application may be needed; tissue samples should be used to determine later usages.
<u>Cotton</u>	1.8 t/ha	Apply between the first true leaf and pinhead square.
	2.25 to 4.5 t/ha	Apply from first bloom to boll fill. Later application may be needed; tissue samples should be used to determine the necessity of applying the product.
<u>Peanuts</u>	2.25 to 4.5 t/ha	Use MAX-IN® ZINC during the vegetative stage in a minimum of 38 litres of water by ground. A second application may be made at pegging.
<u>Potatoes</u>	2.25 to 4.5 t/ha	Apply from emergence to tuber initiation. Later application may be needed; tissue samples should be used to determine later usages.
<u>Soybeans</u>	2.25 to 4.5 t/ha	Apply when soybeans are in the vegetative stage or earlier reproductive stage. Later application may be needed; tissue samples should be used to determine later usages.
<u>Sugar Beets</u>	2.25 to 4.5 t/ha	Apply between the second and 10 true leaf stages. Later application may be needed; tissue samples should be used to determine later usages.
<u>Non-vine vegetable crops</u> Including Edible Beans, Dry Beans, Peppers, Potatoes, Tomatoes and other Fruits	2.25 t/ha	Apply MAX-IN® ZINC starting at pre-bloom or early bloom. Repeat applications may be needed every 10 to 14 days or as tissue sample deem necessary

Crop	Application rate	Remarks
<u>Cucurbits and other vine crops</u> Including Cantaloupes, Watermelons and Honey Dew Melons	2.25 ℓ/ha	Apply MAX-IN® ZINC when runners are 10 to 15 cm in length. May be repeated in 14 to 21 days for a maximum four (4) applications.
<u>Tree Crops</u>	2.25 to 4.5 ℓ/ha	<u>All Nut and Fruit Crops:</u> Apply as recommended by crop advisor. Can be applied in 450 to 1800 litres water per hectare.
<u>Other Crops</u>	-	Apply according to directions of your regional Villa Crop Protection Plant Nutrition Specialist or local agricultural professional, to crops where the addition of zinc would be beneficial.
<u>Turfgrass</u> Golf Courses, Sports Fields, Lawns, and other Turf Areas	10 to 60 ml/100m ²	Apply 10 to 60 ml per 100m ² in 4 to 12 litres of water during the growing season as needed. Apply this product only to your lawn/garden and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden. Do not apply if heavy rain is expected. <u>For Turfgrass Applications:</u> Do not apply near water, storm drains, or drainage ditches.

Aerial Application

- Above use rates should be used in a minimum of 30 litres of water by air.
- Repeat applications 10 to 21 days apart may be necessary.
- A number of variables influence the rate of application necessary to prevent or correct a specific deficiency.
- For best results, follow good agronomic practices and tissue test.
- For specific recommendations for crops in your area, consult your local agricultural professional.

MAX-IN® ZINC is a registered trademark of / is 'n geregistreerde handelsmerk van
WINFIELD SOLUTIONS REGISTRATION HOLDINGS (PTY) LTD.