



novagib[®] 10 SL

A GROWTH REGULATOR FOR THE REDUCTION
OF RUSSETING IN APPLES.

I: 13/1/2017 – Jan2017

'N GROEIREGULEERDER VIR DIE BEHEER VAN
KELK-END VERRUWING IN APPELS.

REG. NO. L 10187 ACT/WET NO. 36 OF/VAN 1947

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:
GIBBERELLIN ACIDS A4 & A7 / GIBBERELLIENSURE A4 & A7 10 G/l

UN NUMBER: NOT REGULATED

MANUFACTURED BY /
VERVAARDIG DEUR:

FINE AGROCHEMICALS LIMITED
HILL END HOUSE WHITTINGTON
WORCESTER WR5 2RQ UK
ENQUIRE@FINE.EU
WWW.FINE.EU

fine

DISTRIBUTED BY /
VERSPREI DEUR:



REGISTRATION HOLDER/REGISTRASIEHOUER
UNIVERSAL CROP PROTECTION (PTY) LTD.
CO, REG. NO. 1983/008184/07
MPY, REG. NR. P.O.BOX / POSBUS 801
KEMPTON PARK, 1620
TEL. (011) 396 2233
WEBSITE / WEBBLAD: WWW.VILLACROP.CO.ZA

BATCH NUMBER /
LOTNOMMER:

DATE FORMULATED /
FORMULERINGSDATUM:

NET VOLUME:
NETTO VOLUME:

1e

NOV/SA/11/17

© NOVAGIB IS A REGISTERED TRADEMARK OF / IS 'N GEREGISTREERDE HANDELSMERK VAN FINE AGROCHEMICALS LTD.



CAUTION
VERSIGTIG



Vir volledige besonderhede, sien ingeslote pamflet.
For full particulars, see enclosed leaflet.

NOVAGIB® 10 SL

Reg. No. L 10187 Act/Wet No. 36 of/van 1947

ACTIVE INGREDIENT/ AKTIEWE BESTANDDEEL:

gibberellin acids A4 & A7 / gibberelliensure A4 & A7 10 g/l

Registration holder / Registrasiehouer:

UNIVERSAL CROP PROTECTION. (PTY) LTD.

Co. Reg. No. 1983/008184/07 Mpy. Reg. Nr.

P.O. Box / Posbus 801, KEMPTON PARK, 1620, Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS**

- Handle the product with care.
- May cause mild skin irritation and severe eye irritation.
- Harmful when swallowed.
- Store in a cool, dry, well-ventilated place.
- Store away from food, feeds, seed, fertilizers and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Do not inhale the spray mist.
- Avoid contact with skin and eyes.
- Wash with soap and water after use and after accidental skin contact.
- In case of eye contact, immediately flush the eyes with clean, cold water for 15 to 20 minutes, holding the eyelid(s) open. If irritation persists, seek medical advice.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face and change of clothing.
- Do not allow drift of spray on to other crops, grazing, rivers, dams or areas not under treatment.
- Clean the applicator and dipping bath after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE** empty containers in the following manner: Invert empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse empty container three times with a volume of clean water equal to one third of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening and dispose of it in a safe way.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

USE RESTRICTIONS

- Delay application when conditions at full bloom are unfavourable. Commence with application when conditions will promote maximum absorption of **NOVAGIB® 10 SL** (this include high relative humidity, cool weather, damp and slow drying conditions at time of spraying or shortly thereafter). Apply in the early morning or evening and avoid rapid drying conditions.
- In the case of rainy weather, delay **NOVAGIB® 10 SL** treatment until no surface droplets are visible on flower parts / fruitlets and leaves. Only apply when a 6-hour period without rainfall is expected. Should

rain falls within 6 hours after application, suboptimal results can be expected. However, don't re-spray in the event of rain falling within 6 hours after treatment.

- Avoid application when temperatures are expected to be lower than 7°C, or exceed 32°C at the time of application.
- Do not spray when wind conditions will promote excessive spray drift.

DIRECTIONS FOR USE: Use only as directed.

- Environmental factors may influence the efficacy of **Gibberellin acid**.
- Use open containers within the season that they were opened.

Compatibility:

Do not apply **NOVAGIB® 10 SL** in tank mixtures with insecticides, foliar feeds or other plant growth regulators.

Mixing Instructions:

- Half fill the spray tank or dipping bath with clean water.
- Add the required amount of **NOVAGIB® 10 SL** to the water while stirring.
- Fill the spray tank/dipping bath with water to the required level, while maintaining agitation, to ensure thorough mixing.
- Maintain agitation during application.
- Apply **NOVAGIB® 10 SL** the same day that the spray mixture was prepared.

Application instructions:

- Use a properly calibrated orchard sprayer that ensures uniform and complete coverage of all fruitlets and foliage and blossoms.
- Apply in a sufficient volume of water to ensure thorough coverage without excessive run-off.
- Determine the spray volume required per hectare using the Tree Row Volume formula. Adjust spray volumes based on tree size and spacing. Spray volumes of approximately 750 to 1000 litres per hectare will be adequate in most cases.

APPLICATION RATES

<i>Crop / Use</i>	<i>Dosage Rate</i>	<i>Remarks</i>
Apples Reduction of calyx end (retiform) russetting in Golden Delicious apples	100 ml / 100 ℓ water	Apply treatments when conditions are likely to promote a high incidence of russetting (e.g. fluctuating weather conditions or orchards with a known severe russet problem). Direct 85 % of the NOVAGIB® 10 SL spray mix as a uniform fine mist spray to the upper 2/3 of the trees. Apply 4 to 5 treatments, commencing at full bloom. Repeat at 7 to 10 day intervals.

NOVAGIB® is a registered trade mark of / is 'n geregisteerde handelsmerk van **Fine Agrochemicals Limited**

PRODUCT WARRANTY CONDITIONS AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications indicated on this label. FINE disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance or other unintended consequences may result because of factors such as abnormal weather conditions, presence of other materials, the manner of application, use of the Product other than in strict accordance with this label's instructions or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, the Buyer shall assume all such risks.

FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product. FINE disclaims any liability whatsoever for special, incidental or consequential damages resulting from the handling or use of this Product.

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ, United Kingdom.