

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY!

Insecticide



T-REX

Reg. No. L 9537 Act/Wet No. 36 of/van 1947

1: 31/3/2014 – June2017

A bait in pellet form for control of Snails, Slugs and Millipedes in agricultural crops and horticulture.

'n Lokaas in korrelvorm vir die beheer van Slakke, Naakslakke en Duisendpote in landbougewasse en tuinbou.

ACTIVE INGREDIENT / AKTIEWE BESTANDEDEL

metaldehyde **30 g/kg** metaldehyd
carbaryl (carbamate) **20 g/kg** karbariel (karbamaat)

IRAC INSECTICIDE GROUP CODE **1A** IRAC INSEKODER GROEPKODE

kg

villa 

Registration holder / Registrasiehouer:

Villa Crop Protection (Pty) Ltd.

Co. Reg. No. / Mpy. Reg. Nr. 1992/002474/07

PO Box / Posbus 10413, Aston Manor, 1630

Tel: 011 396 2233

Website / Webblad: www.villacrop.co.za

UN Number: 3077

Willow Set & Print 011 394-4486



CAUTION
VERSIGTIG



T-REX

Reg. No. L 9537 Act/Wet No. 36 of/van 1947

IRAC INSECTICIDE GROUP CODE / IRAC INSEKODER GROEPKODE: 1A

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

metaldehyde / metaldehyd	30 g/kg
carbaryl (carbamate) / karbariel (karbamaat).....	20 g/kg

Registration holder / Registrasiehouer:

VILLA CROP PROTECTION (PTY) LTD.

Co. Reg. No.: 1992/002474/07 Mpy. Reg. Nr.

PO Box / Posbus 801

KEMPTON PARK, 1620.

Tel. (011) 396 2233

CAUTION / VERSIGTIG**WARNINGS**

- **Withholding periods:**
Do not harvest edible crops within 7 days after application.
- Do not graze or cut for stock food for 7 days after application. Handle with care.
- Toxic if swallowed or inhaled.
- Toxic to fish and highly toxic to birds, mammals and honey bees exposed to direct treatment or residues on blooming shrubs, flowers, weeds and trees. Do not apply this product or allow it to be placed under blooming shrubs, flowers, weeds or trees if bees are visiting the treated area.
- Store the product inside the original container/bag and always keep the container/bag tightly closed.
- Store in a cool, dry, well-ventilated place, away from food, feeds, seed and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.

PROTECTION OF LIVESTOCK

- After filling applicator, clean up spilled pellets so that they are not eaten by animals and birds.
- Dogs find this bait attractive and if ingested it may kill them.
- Do not heap as heaping is unnecessary and such heaps could be eaten by dogs.
- If a dog eats any pellets take it to a veterinarian immediately.
- Do not give dogs Ipecac Syrup.
- Do not treat areas on which poultry feeds.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Avoid contact with the skin and eyes.
- Wear rubber gloves when applying the product.
- Do not eat, drink or smoke whilst preparing or applying or before washing hands and face or change of clothing.
- Do not apply near water bodies such as fishponds.
- Container/bags must be completely emptied before being disposed of. Shake out thoroughly and destroy the empty container/bag thereafter and bury it. **Do not burn the empty container/bag.**
- **Never** re-use the empty container/bag for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

DIRECTIONS FOR USE: Use only as directed.**Use restrictions:**

- Do not apply to edible plant parts or crops where baits may be collected with harvested commodities.
- Do not heap baits.
- To prevent accidental, but fatal poisoning, product should be kept away from ducks, poultry, animals and domestic pets. Contact a veterinary surgeon if pets or livestock are poisoned.

General information:

- **T-REX** kills Snails and Slugs by irreversibly destroying the mucus production cells that are essential for movement as well as by interference with the cholinergic nervous system.
- Under high infestation pressure it may be necessary to repeat application.
- Best results are achieved by application during mild damp weather when Snails and Slugs are most active, but when rain is not imminent.
- For seedlings, apply 5 to 10 days prior to planting.

APPLICATION RATES

<i>Situation</i>	<i>Application rate</i>	<i>Remarks</i>
Apply bait when crops are most vulnerable and when the pest is evident and active. Generally start when rain/moisture triggers Snail and Slug activity, but before significant egg laying occur.		
<u>Row Crops, Orchards and Vines</u> Brown snail <i>(Helix aspersa)</i> White dune snail <i>(Theba pisana)</i> Millipedes (Class: Diplopoda)	5 to 10 kg / ha	Apply as a broadcast application evenly over the ground surface where Snails, Slugs and/or millipedes are active. In grapevines, also apply close to supportive struts. Under normal climatic conditions and infestation pressures, the lower dosage rate should control Brown garden snails effectively. Under certain (not yet quantified) conditions, control of White dune snails may be variable, despite application of the higher dosage rate. Repeat application if infestation re-appears.
<u>Horticulture & Gardens</u> Brown snail <i>(Helix aspersa)</i>	5 to 8 g per 10 m ² area OR 35 to 40 pellets per m ²	Scatter the T-REX bait pellets evenly in flower beds, under dense foliage, shrubs and trees. In case of container plants, place bait pellets close to base of container.