



KEEP OUT OF REACH OF CHILDREN AND ANIMALS. SHAKE BOTTLE WELL BEFORE USE

novagib[®] 10 SL

3: 18/8/2022 – Oct2022

A GROWTH REGULATOR FOR THE REDUCTION
OF RUSSETING IN APPLES.

REG. NO. L 10187 NO. 36 OF 1947

ACTIVE INGREDIENT
GIBBERELLIN ACIDS A4 & A7 10 G/L

UN NUMBER: NOT REGULATED

NO HAZARDS FLAGGED ACCORDING TO GHS CLASSIFICATION

MANUFACTURED BY /
FINE AGROCHEMICALS LIMITED
HILL END HOUSE WHITTINGTON
WORCESTER WR5 2RQ UK
ENQUIRE@FINE.EU
WWW.FINE.EU

fine

© NOVAGIB IS A REGISTERED TRADEMARK OF FINE AGROCHEMICALS LTD.

DISTRIBUTED BY /



REGISTRATION HOLDER
UNIVERSAL CROP PROTECTION (PTY) LTD.
CO. REG. NO. 1983/008184/07
P.O. BOX 801
KEMPTON PARK, 1620
TEL. (011) 396 2233
WEBSITE: WWW.VILLACROP.CO.ZA

BATCH NUMBER /

DATE FORMULATED /

NET VOLUME:

1e

NOV/ZA/1L/22

For full particulars, see enclosed leaflet.

NOVAGIB® 10 SL

Reg. No. L 10187 Act No. 36 of 1947

ACTIVE INGREDIENT:

gibberellin acids A4 & A7 10 g/l

Registration holder:

UNIVERSAL CROP PROTECTION. (PTY) LTD.

Co. Reg. No. 1983/008184/07

P.O. Box 801, KEMPTON PARK, 1620, Tel. (011) 396 2233

WARNINGS

- Handle the product with care.
- Store in a cool, dry, well-ventilated place.
- Store away from food, feeds, seed, fertilizers and other agricultural remedies.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label, as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face and change of clothing.
- Do not allow drift of spray on to other crops, grazing, rivers, dams or areas not under treatment.
- Clean the applicator and dipping bath after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- **TRIPLE RINSE THE EMPTY CONTAINER AS FOLLOWS**: Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three (3) times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

FIRST AID TREATMENT

Inhalation: Remove person from contaminated area to fresh air and assist breathing as needed. Seek medical attention if irritation occurs.

Skin: Remove contaminated clothing and wash before re-using. Flush skin with water. Seek medical attention if skin irritation persists.

Eyes: Flush eyes with clean water. Lift eyelids to facilitate irrigation. If present, remove contact lenses and continue rinsing. Seek medical attention if irritation occurs.

Ingestion: Rinse mouth. Seek medical attention if person feels unwell.

USE RESTRICTIONS

- Delay application when conditions at full bloom are unfavourable. Commence with application when conditions will promote maximum absorption of **NOVAGIB® 10 SL** (this includes high relative humidity, cool weather, damp and slow drying conditions at time of spraying or shortly thereafter). Apply in the early morning or evening and avoid rapid drying conditions.
- In the case of rainy weather, delay **NOVAGIB® 10 SL** treatment until no surface droplets are visible on flower parts / fruitlets and leaves. Only apply when a 6-hour period without rainfall is expected. Should

rain falls within 6 hours after application, suboptimal results can be expected. However, don't re-spray in the event of rain falling within 6 hours after treatment.

- Avoid application when temperatures are expected to be lower than 7°C or exceed 32°C at the time of application.
- Do not spray when wind conditions will promote excessive spray drift.

DIRECTIONS FOR USE: Use only as directed.

- Environmental factors may influence the efficacy of **Gibberellin acid**.
- Use open containers within the season that they were opened.

Compatibility:

Do not apply **NOVAGIB® 10 SL** in tank mixtures with insecticides, foliar feeds or other plant growth regulators.

Mixing Instructions:

- Half fill the spray tank or dipping bath with clean water.
- Add the required amount of **NOVAGIB® 10 SL** to the water while stirring.
- Fill the spray tank/dipping bath with water to the required level, while maintaining agitation, to ensure thorough mixing.
- Maintain agitation during application.
- Apply **NOVAGIB® 10 SL** the same day that the spray mixture was prepared.

Application instructions:

- Use a properly calibrated orchard sprayer that ensures uniform and complete coverage of all fruitlets and foliage and blossoms.
- Apply in a sufficient volume of water to ensure thorough coverage without excessive run-off.
- Determine the spray volume required per hectare using the Tree Row Volume formula. Adjust spray volumes based on tree size and spacing. Spray volumes of approximately 750 to 1000 litres per hectare will be adequate in most cases.

APPLICATION RATES

<i>Crop / Use</i>	<i>Dosage Rate</i>	<i>Remarks</i>
Apples Reduction of calyx end (retiform) russetting in Golden Delicious apples	100 ml / 100 l water	Apply treatments when conditions are likely to promote a high incidence of russetting (e.g., fluctuating weather conditions or orchards with a known severe russet problem). Direct 85 % of the NOVAGIB® 10 SL spray mix as a uniform fine mist spray to the upper 2/3 of the trees. Apply 4 to 5 treatments, commencing at full bloom. Repeat at 7-to-10-day intervals.

NOVAGIB® is a registered trademark of **Fine Agrochemicals Limited**

PRODUCT WARRANTY CONDITIONS AND LIMITATION OF LIABILITY

Fine Agrochemicals Limited ("FINE") warrants that this Product conforms to the specifications indicated on this label. FINE disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for a particular purpose.

It is impossible to eliminate all risks associated with this Product. Plant injury, lack of performance or other unintended consequences may result because of factors such as abnormal weather conditions, presence of other materials, the manner of application, use of the Product other than in strict accordance with this label's instructions or other factors, all of which are beyond the control of FINE or the seller. To the extent consistent with applicable law, the Buyer shall assume all such risks.

FINE's liability under this label shall be limited to the amount of the purchase price or, at the election of FINE, the free replacement of the Product. FINE disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the handling or use of this Product.

© Fine Agrochemicals Limited, Hill End House, Whittington, Worcester, WR5 2RQ, United Kingdom.